CERN-EP-2025-201
2025/10/02

CMS-TOP-24-001

Measurement of the dineutrino system kinematic variables in dileptonic top quark pair production in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

Differential top quark pair production cross sections are measured in the dilepton final states e^+e^- , $\mu^+\mu^-$, and $e^\pm\mu^\mp$, as a function of kinematic variables of the two-neutrino system: the transverse momentum p_T^{vv} of the dineutrino system, the minimum distance in azimuthal angle between $\vec{p}_T^{vv}$ and leptons, and in two dimensions in bins of both observables. The measurements are performed using CERN LHC proton-proton collisions at $\sqrt{s} = 13$ TeV, recorded by the CMS detector between 2016 and 2018, corresponding to an integrated luminosity of 138 fb^{-1} . The measured cross sections are unfolded to the particle level using an unregularized least squares method. Results are compared with predictions by the standard model of particle physics, and found to be in agreement with theoretical calculations as well as Monte Carlo simulations.

Submitted to the Journal of High Energy Physics

1 Introduction

Precision measurements of the top quark pair ($t\bar{t}$) production cross section provide stringent tests for the validity of the standard model of particle physics (SM) while playing a crucial role in the search for new phenomena. The large $t\bar{t}$ data samples collected at the CERN LHC have made several measurements of differential cross sections possible in various $t\bar{t}$ decay channels and at different center-of-mass energies [1–23]. All measurements to date were performed as functions of observables that characterize the kinematic properties of either the visible part of the event, such as jets or charged leptons, or intermediate particles, such as the top (anti)quark or the W boson. However, there are beyond-the-SM (BSM) physics scenarios that mostly modify the invisible part of the signature. Precise and direct measurements of undetected particles in the event, can therefore reveal signs of new phenomena.

This paper presents the first measurements of the differential $t\bar{t}$ cross section as a function of the transverse momentum of the dineutrino system, $p_T^{\nu\nu}$, and the minimum azimuthal distance between $\vec{p}_T^{\nu\nu}$ and the transverse momentum of a charged lepton, $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$. In addition, a two-dimensional (2D) measurement as a function of both observables is performed. For the SM $t\bar{t}$ production process, neutrinos from the leptonic decays of the W boson account for most of the missing transverse momentum $\vec{p}_T^{\text{miss}}$ of the event, while for BSM processes, contributions to $\vec{p}_T^{\text{miss}}$ from undetected particles are also present in the event. These additional contributions weaken the angular correlation between $\vec{p}_T^{\nu\nu}$ and the lepton direction, leading to a broader $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$ distribution. The selection of the two observables for the cross section measurements in this analysis is, therefore, driven by the distinction between the SM $t\bar{t}$ process and potential BSM scenarios that exhibit a comparable signature but include additional sources of undetected particles.

Squarks are superpartners of quarks as predicted by supersymmetry [24]. In Fig. 1, a diagram of $t\bar{t}$ production in the dilepton final state is compared to that of a top squark pair production, as an example of a BSM process [25]. The presence of neutralinos, that neither decay to SM particles nor are detectable within the detector, modifies the kinematic properties of the invisible part of the BSM event that otherwise exhibits a final state identical to that of the $t\bar{t}$ decay. Dedicated searches for top squark pair production in the dilepton final state found significant contributions of $t\bar{t}$ events in the search phase space [26, 27].




Figure 1: In the left diagram, the SM $t\bar{t}$ production process is sketched, whereas the right diagram shows the production of a hypothetical top squark pair, $\tilde{t}_1\tilde{t}_1^*$, with both top squarks decaying to a top quark and a neutralino, $\tilde{\chi}_1^0$. This analysis focuses on signatures, where both of the W bosons decay leptonically.

A precise measurement of the dineutrino system in dileptonic $t\bar{t}$ events requires a good resolution of $\vec{p}_T^{\text{miss}}$, which serves as a proxy for the transverse momentum of the dineutrino system. The magnitude and direction of $\vec{p}_T^{\text{miss}}$ are evaluated from the negative $\vec{p}_T$ sum of all recon-

structed objects in an event, exploiting momentum conservation. The $\vec{p}_T^{\text{miss}}$ resolution is affected by mismeasurements of other particles that can happen due to detector effects and/or as a result of additional proton-proton (pp) collisions in the same or adjacent bunch crossing (pileup). A dedicated correction based on a regression using machine learning methods is developed to improve the $\vec{p}_T^{\text{miss}}$ resolution.

Absolute and normalized differential cross sections are measured in the dilepton final states e^+e^- , $\mu^+\mu^-$, and $e^\pm\mu^\mp$, derived for a fiducial phase space close to the acceptance of the detector and extracted by an unregularized unfolding using a least squares minimization. The results are compared to predictions from Monte Carlo (MC) event generators at the matrix element (ME) level with next-to-leading order (NLO) accuracy in quantum chromodynamics (QCD), which are interfaced with a parton shower (PS) simulation. In addition, the results are compared to fixed-order predictions at NLO and next-to-NLO (NNLO) precision [28].

The paper is structured as follows: Section 2 briefly describes the main features of the CMS detector. In Section 3, the recorded data samples are briefly introduced and settings for MC simulation of signal and background processes are detailed. Section 4 presents the object reconstruction and the event selection, followed by the description of the correction of $\vec{p}_T^{\text{miss}}$ in Section 5. The systematic uncertainties and the results, including comparisons to different predictions, are discussed in Sections 6 and 7, respectively. Finally, the analysis is summarised in Section 8. Tabulated results can be found in a HEPData record [29].

2 The CMS detector

The CMS apparatus [30, 31] is a multipurpose, nearly hermetic detector, designed to trigger on [32–34] and identify electrons, muons, photons, charged and neutral hadrons [35–37]. Its central feature is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are reconstructed using gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of 4 μ s [32]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [33].

More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [30, 31].

3 Data and simulated samples

The measurements presented in this paper are based on pp collision data recorded between 2016 and 2018 at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 138 fb⁻¹ [38–40] with 36.3, 41.5, and 59.8 fb⁻¹ of data recorded in 2016, 2017, and 2018, respectively.

Both signal and the SM background processes are simulated using MC event generators. Simulated samples are produced separately for each data-taking period to match the corresponding

pileup and detector conditions. All simulated samples use parton distribution functions (PDFs) as implemented in NNPDF3.1 [41, 42] at NNLO accuracy in perturbative QCD. All ME event generators are interfaced with PYTHIA v8.230 [43] for the simulation of the parton showering and hadronization using the CP5 tune [44] unless stated otherwise in the text. The detector response is simulated with GEANT4 [45].

The $t\bar{t}$ production process is simulated using the POWHEG v2 [46–48] event generator at NLO precision with the top quark mass set to 172.5 GeV. Top quark pairs in which both W bosons decay into either an electron or muon belong to the signal sample and are referred to as “ $t\bar{t} (\ell\ell)$ ” and/or “dileptonic $t\bar{t}$ ” in the following. All other decay channels, including the production of one or more tau leptons, are denoted as “ $t\bar{t}$ other”. Additional $t\bar{t}$ samples using the same event generator are produced with modified parameters for the underlying event tune, color reconnection scheme, jet matching scheme, and top quark mass, and are used in the estimation of the impact of theoretical uncertainties. A $t\bar{t}$ sample generated with MADGRAPH5_aMC@NLO v2.4.2 (MG5) [49] at NLO accuracy and interfaced with PYTHIA through the FxFx [50] matching and merging scheme is used for the determination of corrections that improve the experimental $\vec{p}_T^{\text{miss}}$ resolution detailed in Section 5. This sample is referred to as MC@NLO+PYTHIA in the following. Another alternative $t\bar{t}$ sample is generated with the POWHEG v2 generator interfaced with HERWIG v7.1 [51] using the CH3 tune [52], and is referred to as POWHEG+HERWIG.

Background processes are modelled using different MC event generators. Drell–Yan events with additional jets and two leptons in the final state (DY+jets) are produced at NLO using MG5. Events are passed through PS and hadronization using the FxFx merging scheme.

The same event generator is used for the production of a W boson that subsequently decays to a lepton and neutrino. Events contain up to four additional jets, modelled at leading order in perturbative QCD, and matched to jets produced during PS and hadronization with the MLM scheme [53]. The electroweak top quark productions, referred to as “single top” hereafter, are modelled at NLO accuracy using POWHEG v2 [54] for the tW process and the t -channel production, and using MG5 in the case of the s -channel production. For the tW process, the description based on the diagram subtraction (DS) scheme is applied [55] in order to handle the overlap and interference with $t\bar{t}$ production. Another tW sample based on the diagram removal (DR) scheme [55] is used for the estimation of theoretical uncertainties.

The production of $t\bar{t}$ in association with a W or Z boson ($t\bar{t}W/Z$) is modeled at NLO accuracy using MG5. Diboson production processes, including WW, WZ, and ZZ, are generated at leading order using PYTHIA.

The expected yield of $t\bar{t}$ events is determined using a cross section calculated with the TOP++ [56] program at NNLO plus next-to-next-to-leading-logarithmic accuracy [57–60], assuming a top quark mass of 172.5 GeV. The simulated p_T spectra of top quarks are reweighted to NNLO cross sections using NNLO/NLO K-factors [61]. Yields from the W+jets and high-mass DY+jets processes are calculated using NNLO accuracy cross sections [62, 63], while the yield of low-mass DY+jets and the $t\bar{t}W/Z$ processes are calculated using NLO accuracy cross sections from the MG5 generator. A cross section calculation at approximate NNLO is used for the single top tW [64] process, while yields from the single top s - and t -channel processes are determined using NLO cross sections [65, 66]. Cross sections calculated at NNLO accuracy [67] are used to determine the WW yield, whereas NLO cross sections [68] are applied to determine the WZ and ZZ diboson process yields.

4 Object reconstruction and event selection

Events passing dilepton and single-lepton triggers are used for the offline analysis. The trigger criteria vary by data-taking period. For dielectron and electron-muon triggers, the transverse momentum p_T of the (sub)leading lepton is required to pass a threshold of 23 (12) GeV. The maximum p_T threshold on the (sub)leading muon in dimuon triggers is 17 (8) GeV. The threshold on the lepton p_T is 27 (32) GeV in 2016 (2017–2018) for single electron triggers, and 24 (27) GeV in 2016/2018 (2017) for single muon triggers.

The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [69]. The particle-flow (PF) algorithm [70] aims to reconstruct and identify each individual particle in the event, with an optimized combination of information from various elements of the CMS detector. The energy of photons is obtained from the ECAL measurement. The energy of electrons is determined from a combination of the electron momentum at the PV, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The energy of muons is obtained from the curvature of the corresponding track. The energy of charged hadrons is determined from a combination of their momenta measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

Electron candidates are vetoed if their associated ECAL cluster is reconstructed within the transition region of the ECAL, which corresponds to the η range of $1.44 < |\eta| < 1.57$. Misidentified and nonprompt electrons are further suppressed by implementing selections based on discriminating quantities, such as shower shape variables, relative isolation, longitudinal and transverse impact parameters with respect to the PV, and photon conversion veto observables. For electrons, the relative isolation is defined as the p_T sum of all neutral and charged hadrons, and photon candidates within an angular distance $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ of 0.3 from the electron in the η - ϕ space, divided by the p_T of the electron candidate. Depending on the p_T and η of the electron candidate, a maximum relative isolation in the range 0.05–0.1 is enforced. Muon candidates are required to have matching tracks in the inner tracker and the muon system, with track quality criteria, relative isolation, and impact parameter requirements suppressing nonprompt or misidentified muons. For muons, the relative isolation is based on contributions within a distance of 0.4 from the muon candidate, applying a maximal relative isolation of 0.15. For both electrons and muons, the relative isolation is corrected for pileup effects by removing charged PF candidates associated with pileup vertices, and by estimating the contribution of neutral particles from pileup to be subtracted from the p_T sum [71, 72].

Jets are clustered using the anti- k_T algorithm [73, 74] with a distance parameter of 0.4. Jets within an angular distance $\Delta R < 0.4$ of selected leptons are not considered in further steps of the analysis. To reduce contributions from jets originating from pileup vertices, additional requirements for jets reconstructed with $p_T < 50$ GeV are applied [75]. Jets originating from the hadronization of b quarks are identified with the DEEPJET algorithm [76], using a working point corresponding to an efficiency of up to 92% for b quark jets and a misidentification rate of 10 (40–50)% for jets originating from light-flavor quarks and gluons (c quarks).

The missing transverse momentum vector $\vec{p}_T^{\text{miss}}$ is computed as the negative vector sum of the transverse momenta of all the PF candidates in an event, and its magnitude is denoted as p_T^{miss} [77]. The $\vec{p}_T^{\text{miss}}$ is modified to account for corrections to the energy scale of the reconstructed jets in the event and denoted as $\vec{p}_{T,\text{PF}}^{\text{miss}}$. The pileup-per-particle identification (PUPPI)

algorithm [78] is applied to reduce the pileup dependence of the $\vec{p}_T^{\text{miss}}$ observable. The $\vec{p}_T^{\text{miss}}$ computed from the PF candidates weighted by their probability to originate from the PV [77] is denoted as $\vec{p}_{T,\text{PUPPI}}^{\text{miss}}$. Further corrections to $\vec{p}_T^{\text{miss}}$ are derived in this analysis using a deep neural network (DNN) in order to improve the p_T^{miss} resolution in dileptonic $t\bar{t}$ events, especially at high values of p_T^{miss} . This procedure is further detailed in Section 5 and the corrected $\vec{p}_T^{\text{miss}}$ is denoted as $\vec{p}_{T,\text{DNN}}^{\text{miss}}$. Anomalous high- p_T^{miss} events can occur due to a variety of failures in the object reconstruction, detector malfunctions, and/or noncollision backgrounds. Such events are rejected by event filters designed to identify more than 85–90% of the spurious high- p_T^{miss} events with a misidentification rate of less than 0.1% [77].

The event selection closely follows that of Ref. [18] with the exception that the requirement on missing transverse momentum is applied to $\vec{p}_{T,\text{DNN}}^{\text{miss}}$ rather than $\vec{p}_{T,\text{PF}}^{\text{miss}}$. Events with two oppositely charged leptons that fulfill $|\eta| < 2.4$ and $p_T > 25$ (20) GeV for the leading (subleading) lepton are selected and split into the same-flavor $e^+e^-/\mu^+\mu^-$ and the different-flavor $e^\pm\mu^\mp$ channels. Events containing additional leptons (electrons or muons) with $p_T > 15$ GeV are rejected. Contributions from low-mass resonances decaying to two leptons are suppressed by requiring a minimal invariant mass $m_{\ell\ell}$ of the two leptons of 20 GeV. In order to reduce the contribution from DY+jet events, the events in the same-flavor channels are required to fulfill $p_{T,\text{DNN}}^{\text{miss}} > 40$ GeV and are vetoed if $m_{\ell\ell}$ is compatible with the Z boson mass [79] within 15 GeV. Selected events must contain at least two reconstructed jets with $p_T > 30$ GeV and $|\eta| < 2.4$, of which at least one is b tagged [80].

The numbers of expected events from simulation and of observed events in data are shown in Table 1. The signal process $t\bar{t}$ ($\ell\ell$) comprises 78% of the selected events. The largest background contribution arises from other $t\bar{t}$ processes, contributing to 13% of the selected sample. It mostly contains events where at least one W boson decays to a τ lepton. The next background contributions in size are single top and DY processes, with 4.2 and 4.0% contributions to the selected samples, respectively. Distributions of a selected set of kinematic observables are shown in Fig. 2. The simulated samples reproduce the distributions in data well.

Table 1: Data and MC simulation yields after the event selection, combined for all data-taking periods and split by channels. The uncertainties on the expected yields include systematic and statistical uncertainties (discussed in Section 6). The relative contribution in percent of each process to the total expected yield of a channel is given in parentheses.

Process	Yields / 10^3			
	e^+e^-	$\mu^+\mu^-$	$e^\pm\mu^\mp$	All
$t\bar{t}$ ($\ell\ell$)	116 (75.2%)	221 (74.1%)	531 (80.2%)	867 (77.9%)
$t\bar{t}$ other	18.2 (11.8%)	38.6 (12.9%)	90.5 (13.7%)	147 (13.2%)
Single top	6.15 (4.0%)	11.7 (3.9%)	28.4 (4.3%)	46.3 (4.2%)
DY+jets	12.6 (8.2%)	24.8 (8.3%)	6.87 (1.0%)	44.2 (4.0%)
Diboson	0.70 (0.5%)	1.24 (0.4%)	2.43 (0.4%)	4.36 (0.4%)
$t\bar{t}W/Z$	0.49 (0.3%)	0.89 (0.3%)	1.82 (0.3%)	3.20 (0.3%)
W+jets	0.08 (0.0%)	0.08 (0.0%)	0.88 (0.1%)	1.03 (0.1%)
Sum MC	154 ± 10	298 ± 14	662 ± 27	1114 ± 47
Data	152	302	650	1103




Figure 2: Observed (black markers) and expected (filled histograms) distributions of leading lepton p_T (upper left), leading jet p_T (upper right), the number of jets (lower left), and the number of b tagged jets (lower right), after event selection. The hatched (grey) areas denote the systematic (total) uncertainties on the expected yields. Vertical and horizontal error bars on the data points represent the statistical uncertainty and the bin width, respectively. Events from all data-taking periods and all channels are combined. The lower panel of each plot shows the ratio between observed and expected distributions. The last bin includes all events above the plotted range.

5 Improving the p_T^{miss} resolution for dileptonic $t\bar{t}$ events

Since the $\vec{p}_T^{\text{miss}}$ is used as a measure of the $\vec{p}_T$ of the two prompt neutrinos ($\vec{p}_T^{l\nu}$) produced in dileptonic $t\bar{t}$ decays, it is important to ensure an accurate reconstruction of its magnitude and direction. Poor resolution or large biases of the reconstructed $\vec{p}_T^{\text{miss}}$, denoted as $\vec{p}_{T,\text{rec}}^{\text{miss}}$, can compromise the stability of the unfolding process and lead to a reduced number of bins for the final measurements. Other than those coming from the two neutrinos from top quark decays, there are two additional sources that contribute to the evaluation of $\vec{p}_T^{\text{miss}}$ in dileptonic $t\bar{t}$ events. The first source corresponds to the production of nonprompt neutrinos from semileptonic hadron decays in jets, especially in jets originating from b quarks. These nonprompt neutrinos lead to genuine $\vec{p}_T^{\text{miss}}$ and are therefore included in the calculation of the generated p_T^{miss} of an event. This quantity, defined as the magnitude of the vector sum of all simulated neutrino $\vec{p}_T$ is denoted as $\vec{p}_{T,\text{gen}}^{\text{miss}}$. The second source is the mismeasurement of particle momenta during reconstruction, with the largest impact arising from the mismeasurements in jets. A regression based on a DNN, targeting $\vec{p}_{T,\text{gen}}^{\text{miss}}$, is derived to correct for the mismeasurements. The

method, detailed in the following, corrects for reconstruction effects based on the targeted final state, while being independent of the exact particle kinematic properties and the description of hadronization.

A feed-forward, fully-connected DNN with two output nodes is constructed and trained using the TENSORFLOW [81] backend with a KERAS [82] interface. In order to simultaneously correct the direction and magnitude of $\vec{p}_T^{\text{miss}}$, the two output nodes of the DNN are the x and y components of the difference between $\vec{p}_{T,\text{PUPPI}}^{\text{miss}}$ and $\vec{p}_{T,\text{gen.}}^{\text{miss}}$. The output of the DNN is then used to correct the $\vec{p}_{T,\text{PUPPI}}^{\text{miss}}$. To avoid a potential bias towards the nominal $t\bar{t}$ sample generated with POWHEG+PYTHIA, which is further used in the unfolding procedure, the network is trained based on the alternative MC@NLO+PYTHIA $t\bar{t}$ sample. The training is performed on dileptonic $t\bar{t}$ events passing the selection detailed in Section 4, but omitting the requirement on $p_{T,\text{DNN}}^{\text{miss}}$. The training sample is reweighted so that the $p_{T,\text{gen.}}^{\text{miss}}$ distribution is uniform up to 500 GeV, and a constant weight is applied to events above 500 GeV. This allows the DNN to perform on a wider $\vec{p}_{T,\text{gen.}}^{\text{miss}}$ range rather than being biased to a specific range with the largest population in the nominal training sample. To account for varying data-taking conditions, the training is performed separately for each data-taking period.

The DNN is based on 17 input observables, including the x and y components of $\vec{p}_{T,\text{PUPPI}}^{\text{miss}}$, $\vec{p}_{T,\text{PF}}^{\text{miss}}$, the $\vec{p}_T$ sum of all jets, the $\vec{p}_T$ of the leading and subleading jets, and the $\vec{p}_T$ of the leading lepton. Additional inputs include the invariant mass of the two leptons and all jets, the invariant mass of the two leading jets, the energy of the leading jet, and the scalar p_T sum of all jets. It is confirmed that all inputs are well modeled in the simulation by fitting the distribution of each input observable to data. All systematic uncertainties (discussed in Section 6) are included in the fit as nuisance parameters. In each case, the goodness of fit is evaluated based on a saturated model [83], as implemented in the CMS statistical analysis tool, COMBINE [84]. The procedure is repeated for every 2D distribution of two inputs to ensure a correct description of correlations between inputs by the simulation.

The performance of $p_{T,\text{DNN}}^{\text{miss}}$ is compared to $p_{T,\text{PF}}^{\text{miss}}$ and $p_{T,\text{PUPPI}}^{\text{miss}}$ in Fig. 3, using a simulated sample corresponding to the 2018 data-taking period. The mean (μ) and the resolution (standard deviation, σ) of the difference between $p_{T,\text{gen.}}^{\text{miss}}$ and $p_{T,\text{rec.}}^{\text{miss}}$ is shown as a function of the $p_{T,\text{gen.}}^{\text{miss}}$ and the number of PVs. As a function of both observables, $p_{T,\text{DNN}}^{\text{miss}}$ improves on the resolution, and inclusively it shifts the mean closer to zero in the phase space of this analysis compared to the other p_T^{miss} reconstruction algorithms. Overall, the resolution of p_T^{miss} is improved by approximately 15% by the DNN with respect to $p_{T,\text{PUPPI}}^{\text{miss}}$, while the resolution of $\phi(\vec{p}_T^{\text{miss}})$ is improved by around 12%. The improvements allow for a finer binning in the differential measurements of the target observables as they result in a reduction of bin-to-bin migration while ensuring a stable unfolding.

6 Systematic uncertainties

The differential cross section measurements are affected by various sources of systematic uncertainty. They can be grouped into two categories, namely experimental and theoretical uncertainties. The impact of each source is evaluated by varying the uncertainty source by ± 1 standard deviation or by using alternative simulation settings. These variations can affect the response matrix used in the unfolding procedure, as well as the background estimation. The resulting change in the extracted cross sections is then taken as the uncertainty from the particular systematic variation. Because of different detector and/or simulation conditions, dedicated correlation models between different data-taking periods are employed for each source of sys-




Figure 3: Difference between $p_{T, \text{gen}}^{\text{miss}}$ and $p_{T, \text{rec}}^{\text{miss}}$ as a function of the $p_{T, \text{gen}}^{\text{miss}}$ (upper) and the number of primary vertices (lower) for signal events. The mean difference between the generated and the $p_{T, \text{rec}}^{\text{miss}}$ per bin is shown as solid line, while the dashed line shows the standard deviation σ , which corresponds to the resolution. The results for p_T^{miss} corrected by the DNN regression (thick light blue), derived with the PUPPI algorithm (thin red), and the PF algorithm (thinner orange) are shown. A simulated sample for the 2018 data-taking period is used.

tematic uncertainty. Unless stated otherwise, the uncertainty is assumed to be fully correlated between different years of data taking.

Experimental uncertainties include those originating from jet reconstruction, corrections applied to simulated samples to improve the shape agreement of data and simulation, normalization of background processes, integrated luminosity measurement, and determination of unclustered energy. The impact from the jet energy scale (JES) uncertainty is estimated by shifting all jet momenta according to seven independent uncertainty sources, which have different treatments of correlations between the data-taking periods. The corresponding shifts in the jet momentum are propagated to p_T^{miss} [77, 85]. The uncertainty on the jet energy resolution (JER) is introduced when smearing jet energies in simulation to match the resolution observed in data. To evaluate its impact, the smearing is repeated with the JER shifted within its uncertainty. This uncertainty is assumed to be uncorrelated between the data-taking periods.

Correction factors are applied to simulation to account for the differences with data when describing b tagging, lepton selection, lepton energy scale, identification of jets from pileup, trigger selection, and ECAL first level trigger inefficiencies [71]. Each of these correction factors is varied within its uncertainty to estimate its impact on the measured cross section. The statistical component of the uncertainty related to the muon identification efficiency, as well as trigger efficiencies, is considered to be uncorrelated between the data-taking periods. Variations in lepton energy scale are propagated to p_T^{miss} . In addition, the corrections to b tagging efficiencies are re-evaluated separately for systematic variations in JES, JER, identification of jets from pileup, and most of the theoretical uncertainties.

The uncertainties on the integrated luminosities are 1.2, 2.3, and 2.5%, for 2016, 2017, and 2018, respectively, with partial correlations between data-taking periods. This results in a combined uncertainty of 1.6% [38–40].

The distribution of additional pp collisions in simulation is reweighted according to the expected distribution in data assuming a total inelastic pp cross section of 69.2 mb [86]. The reweighting is repeated shifting the inelastic pp cross section by $\pm 4.6\%$ to derive the corresponding uncertainty.

The normalization of all background processes directly impacts the cross section measurements. The magnitudes of the corresponding uncertainties applied to the normalization are taken from Ref. [87]. The largest background contribution, denoted as $t\bar{t}$ other, is assigned a 5% normalization uncertainty. For single top and DY+jets production, a 20% and a 30% normalization uncertainty is assigned, respectively. Subdominant background processes, such as diboson production, W+jets, and $t\bar{t}W/Z$ production are assumed to have a 30% normalization uncertainty. Each background normalization is varied individually within the given uncertainty. The simulated samples are normalized to the integrated luminosities of the corresponding data-taking periods [38–40]. The sample normalization is shifted according to the integrated luminosity uncertainties, which are split into uncorrelated and correlated components between the different data-taking periods.

The p_T^{miss} measurement is affected by energy deposits that remain unclustered. To estimate their impact, the energy from charged and neutral hadrons, as well as photons and hadrons reconstructed in the η region covered by the forward hadron calorimeter, is shifted according to their energy resolutions and propagated to p_T^{miss} . The resulting uncertainty is assumed to be uncorrelated between data-taking periods.

Theoretical uncertainty sources include variations that affect all simulated samples, as well as variations that are only applied to the $t\bar{t}$ simulation. The impact of initial-state radiation and

final-state radiation is estimated by shifting their scale by a factor of two and a factor of one half, separately for both initial-state and final-state radiation scales. The uncertainty on the choice of renormalization and factorization scales is estimated in a similar way, shifting the scales up and down by a factor of two, both individually and simultaneously. The uncertainty on the cross section results is given by the envelope of the corresponding shift variations in the measurements. Uncertainties due to the matching of ME and PS calculations are taken into account by reweighting the nominal $t\bar{t}$ samples using weights derived with the DNNs using the classification for tuning and reweighting approach [88, 89] to shift the h_{damp} parameter of the POWHEG event generator. This parameter controls the merging between ME and PS and regulates the high p_T radiation. The impact of the choice of the scheme to handle the overlap between tW and $t\bar{t}$ production [55] is estimated by taking the difference between the tW sample based on DR and DS as additional uncertainty into account.

The remaining theoretical uncertainties are only taken into account for the $t\bar{t}$ simulation. Alternative $t\bar{t}$ samples are used to estimate the impact of varying the top quark mass by 1 GeV, using different tunes for the underlying event modeling as described in Ref. [44], employing two alternative approaches for color reconnection, and enabling early resonance decays [90]. The energy measurement of jets from b quarks is impacted by semileptonic decays of b hadrons. The assumed branching fractions, $\mathcal{B}$, taken from Ref. [79], are shifted within the reported uncertainty to estimate the effect on the cross section measurements. The impact of reweighting the top quark p_T spectra to NNLO accuracy is evaluated in the cross section measurement by repeating the measurement without the reweighting and taking the difference as a one-sided uncertainty.

Uncertainties associated with the PDF description are evaluated by reweighting the simulated $t\bar{t}$ sample using weights based on 100 eigenvectors of the NNPDF3.1 Hessian error PDF set [41, 42], keeping the overall normalization fixed. The envelope of all 100 variation shifts is taken as a systematic uncertainty.

The impact of the systematic uncertainties on the cross section measurements is shown in Fig. 4. For the $p_T^{v\nu}$ measurement, JES is the dominant experimental uncertainty for most of the bins. At large $p_T^{v\nu}$, the choice of the tW – $t\bar{t}$ overlap removal scheme is the largest uncertainty contribution, followed by the normalization of the single top production background and the ME scale, which also mainly affects the single top background estimation. For the $\min[\Delta\phi(\vec{p}_T^{v\nu}, \vec{p}_T^\ell)]$ measurement, the JES uncertainty is dominant at low values, while at large distances the uncertainty related to the lepton reconstruction has the largest contribution. The 2D measurement shows similar behavior to the $p_T^{v\nu}$ measurement, i.e., for most bins the JES uncertainty is the largest systematic uncertainty. At large $p_T^{v\nu}$ in the lowest $\min[\Delta\phi(\vec{p}_T^{v\nu}, \vec{p}_T^\ell)]$ bin, the choice of the tW – $t\bar{t}$ overlap removal scheme is the dominant uncertainty. For the bin with the highest $p_T^{v\nu}$ and $\min[\Delta\phi(\vec{p}_T^{v\nu}, \vec{p}_T^\ell)]$, sizeable contributions also arise from the ME-PS matching and ME scale uncertainties.

The bin-to-bin correlations between different sources of uncertainties are extracted from the covariance matrices that include separate terms for statistical and systematic contributions. The statistical component, encoding uncertainties after the unfolding, may contain off-diagonal elements originating from correlations introduced in the unfolding procedure. The same matrices are used to calculate χ^2 values when comparing predictions to the measured results. The procedure of constructing the covariance matrices is described in detail in Appendix A of Ref. [87].




Figure 4: Breakdown of the relative uncertainties from experimental (left) and theoretical (right) uncertainties on the differential cross section measurement as a function of p_T^{vv} (upper), $\min[\Delta\phi(\vec{p}_T^{vv}, \vec{p}_T^\ell)]$ (middle), and both variables (lower). In the last case, each group of four bins corresponds to p_T^{vv} bins, and $\min[\Delta\phi(\vec{p}_T^{vv}, \vec{p}_T^\ell)]$ bin edges are indicated by vertical solid lines. The statistical uncertainty (dark grey) takes the size of available event counts in both the simulated samples and the recorded data into account. The last bin includes all events above the plotted range.

7 Results

The detector-level distributions of $p_{T,DNN}^{\text{miss}}$, $\min[\Delta\phi(\vec{p}_{T,DNN}^{\text{miss}}, \vec{p}_T^\ell)]$, and the 2D distribution of both observables are compared to the expectation from simulation in Fig. 5. In general, good agreement between the observed data and the simulation is found. For the 2D distribution, a slight over fluctuation at large $\min[\Delta\phi(\vec{p}_{T,DNN}^{\text{miss}}, \vec{p}_T^\ell)]$ and $p_{T,DNN}^{\text{miss}}$ is found in data, still compatible with simulation considering the large statistical uncertainties in this phase space.

The differential cross sections are derived for a fiducial phase space defined in Table 2, closely following the definitions detailed in Ref. [87], with the additional requirement of $p_T^{vv} > 40$ GeV in the same-flavor channel. The binning applied in the final measurements is optimized taking into account the detector resolution, migration effects, and statistical uncertainties. The differ-




Figure 5: The observed (black markers) and simulated distributions of $p_{T,DNN}^{\text{miss}}$ (upper left), $\min[\Delta\phi(\vec{p}_{T,DNN}^{\text{miss}}, \vec{p}_T^\ell)]$ (upper right), and the 2D distribution (lower) with both observables are shown. Events from all data-taking periods and all channels are combined. The hatched (grey) areas indicate the systematic (total) uncertainty on the simulation. Vertical and horizontal error bars on the data points represent the statistical uncertainty and the bin width, respectively. The lower panel of each plot shows the ratio of observed data to the expectation from MC simulation in each bin. The last bin includes all events above the plotted range.

Table 2: Definition of the fiducial phase space for the same-flavor and the different-flavor channels.

Parameter	$e^+e^-/\mu^+\mu^-$	$e^\pm\mu^\mp$
Lepton p_T	$>20 \text{ GeV}$	
Lepton $ \eta $	<2.4	
Number of leptons	$=2$	
Invariant mass of the two leptons	$>20 \text{ GeV}$	
Jet p_T	$>30 \text{ GeV}$	
Jet $ \eta $	<2.4	
$\Delta R(\text{Jet}, \text{Lepton})$	>0.4	
Number of b jets	≥ 2	
Dineutrino transverse momentum ($p_T^{\nu\nu}$)	$>40 \text{ GeV}$	—

ential cross section results are extracted from the observed distributions in two steps. In the first step, the estimated contributions from all backgrounds except for $t\bar{t}$ other are subtracted from the observed distributions. In the second step, the observed distributions are unfolded to the fiducial phase space. Within the second step, the contributions from $t\bar{t}$ other are removed by applying a fixed signal fraction per bin, derived based on the ratio of $t\bar{t}$ signal events to all selected $t\bar{t}$ events.

For the unfolding procedure, the TUNFOLD algorithm [91] is used, performing a least squares minimization without regularization. The choice of the method is motivated by dedicated unfolding studies, where distorted particle-level distributions as well as pseudodata based on the alternative MC@NLO+PYTHIA $t\bar{t}$ sample were unfolded using the nominal response matrix. The selected method showed the most stable performance in comparison with a simple bin-by-bin (BBB) unfolding where only the diagonal elements of the response matrix are taken into account as well as with respect to a least squares minimization with regularization. In addition, closure tests accounting for potential BSM contributions based on the top squark pair production scenario shown in Fig. 1 were performed, for a top squark mass of 525 GeV and a neutralino mass of 350 GeV. Within these tests, pseudodata based on the sum of the nominal $t\bar{t}$ signal prediction and the prediction for the top squark pair production scenario scaled up by an additional factor of ten were unfolded using the nominal response matrix. The corresponding results for the $p_T^{\nu\nu}$ and the 2D distribution can be found in Fig. 6. Based on the nominal unfolding approach, the expected distributions, where both neutralinos are included in the particle-level definition, can be correctly reproduced. This result confirms that the measurement reproduces deviations from the distribution assumed in the MC simulation, which was used in the construction of the response matrix. The normalized differential cross sections are also derived, with the measured cross section in each bin divided by the total fiducial cross section, allowing for the reduction of several systematic uncertainties.

The measured differential cross sections are compared to five theoretical predictions, namely predictions from the POWHEG+PYTHIA, POWHEG+HERWIG, and MC@NLO+PYTHIA $t\bar{t}$ samples, and fixed-order calculations [28] at NLO and NNLO accuracy in QCD. The fixed-order calculations use the narrow-width approximation for the top quark decays and include ME scale uncertainties and, in case of the NNLO prediction, also PDF uncertainties. For the three MC-based predictions, a normalization uncertainty of 5% as well as ME scale uncertainties are taken into account. For the NLO MC simulations, only the acceptance and efficiency effects for the fiducial volume are considered. More details on the cross section extraction can be found in Ref. [87], which only differs in the choice of regularization.

The differential signal cross sections and their comparisons to the theoretical predictions are shown in Figs. 7–8. The corresponding χ^2 tests for all measurements, both with and without the inclusion of uncertainties on the predictions, are summarized in Table 3, with the corresponding p -values [79] given in Table 4.

In general, all predictions are in agreement with the absolute measurements. In the absolute $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$ measurement, small shape differences between the measured cross section and the five predictions are observed. These differences match observations from previous measurements of the azimuthal angle between two leptons $\Delta\phi(\ell, \ell)$ [18, 92], which is strongly correlated with $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$. Slightly larger χ^2/ndf values (where ndf is the number of degrees of freedom) are observed for the 2D differential cross section measurement. While MC@NLO+PYTHIA overpredicts the cross sections in most of the measured bins for the $p_T^{\nu\nu}$ distribution in comparison with other predictions, its χ^2/ndf is found to be close to 1 due to the correlated bin uncertainties. Including the uncertainties on the predictions reduces the χ^2/ndf




Figure 6: Result of the closure test based on simulation accounting for potential BSM contributions based on a top squark pair production scenario with a stop mass of 525 GeV and a neutralino mass of 350 GeV. The potential BSM contribution is scaled by a factor of ten. The test is performed for the p_T^{VV} (upper) and 2D measurement (lower) using the nominal (black circles), the regularized (orange triangles), and the bin-by-bin unfolding (purple squares), based on all data-taking periods combined. The unfolded distributions are compared to the expected distribution (red solid line) based on the sum of the nominal dileptonic $t\bar{t}$ signal sample and BSM signal with the corresponding $\chi^2 / (\text{number of degrees of freedom})$ values given in the legend in square brackets. The distribution used for the response matrix is shown in blue (dotted line). The error bars correspond to the statistical uncertainty. The lower part shows the ratios between the unfolded and the expected distributions.




Figure 7: The measured differential signal cross sections (black markers) as functions of $p_T^{\nu\nu}$ (upper), $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$ (middle), and double-differential in both observables (lower) are shown. The theoretical predictions from POWHEG+PYTHIA (dark red), POWHEG+HERWIG (orange), MC@NLO+PYTHIA (purple), and the fixed-order NLO (light blue) and NNLO (brown) calculations are compared to the measurement. The cross sections in the last $p_T^{\nu\nu}$ bin of the upper (lower) plot are determined for $p_T^{\nu\nu} > 410$ (200) GeV and normalised to a bin width of 90 (200) GeV for visualisation purposes. The total (statistical) uncertainty on the measurement is shown as an orange (dark grey) band. The lower panel of each plot shows the ratio between theoretical predictions and the measurement.




Figure 8: The measured normalized differential signal cross sections (black markers) as functions of $p_T^{\nu\nu}$ (upper), $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$ (middle), and double-differential in both observables (lower) are shown. The theoretical predictions from POWHEG+PYTHIA (dark red), POWHEG+HERWIG (orange), MC@NLO+PYTHIA (purple), and the fixed-order NLO (light blue) and NNLO (brown) calculations are compared to the measurement. The cross sections in the last $p_T^{\nu\nu}$ bin of the upper (lower) plot are determined for $p_T^{\nu\nu} > 410$ (200) GeV and normalised to a bin width of 90 (200) GeV for visualisation purposes. The total (statistical) uncertainty on the measurement is shown as an orange (dark grey) band. The lower panel of each plot shows the ratio between theoretical predictions and the measurement.

Table 3: Results of the χ^2 tests for the absolute and normalized differential cross section measurements for each of the predictions. The χ^2 values including the uncertainties on the predictions are given in parentheses.

Measurement	ndf	χ^2 (including uncertainties)				
		POWHEG +PYTHIA	POWHEG +HERWIG	MC@NLO +PYTHIA	NLO	NNLO
$p_T^{\nu\nu}$	9	6.8 (1.6)	8.0 (1.0)	9.0 (3.9)	7.1 (1.4)	12.5 (2.5)
$\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$	6	7.3 (1.5)	7.4 (1.4)	11.5 (2.7)	12.3 (2.5)	11.9 (4.0)
2D	12	18.8 (4.9)	22.4 (3.7)	31.6 (8.6)	24.0 (3.5)	15.8 (4.2)
norm. $p_T^{\nu\nu}$	8	6.4 (3.8)	8.3 (3.2)	6.4 (5.8)	6.7 (1.9)	13.3 (5.5)
norm. $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$	5	7.2 (7.0)	7.2 (6.7)	9.2 (8.4)	11.7 (5.4)	11.7 (5.8)
norm. 2D	11	18.1 (16.0)	21.4 (14.5)	28.2 (22.8)	23.1 (7.0)	14.8 (8.5)

Table 4: The p -value of the χ^2 tests for the absolute and normalized cross section measurements for each of the predictions. The resulting p -values including uncertainties on the predictions are given in parentheses.

Measurement	p -value (including uncertainties)				
	POWHEG +PYTHIA	POWHEG +HERWIG	MC@NLO +PYTHIA	NLO	NNLO
$p_T^{\nu\nu}$	0.66 (1.00)	0.53 (1.00)	0.44 (0.92)	0.63 (1.00)	0.19 (0.98)
$\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$	0.29 (0.96)	0.28 (0.97)	0.07 (0.84)	0.06 (0.87)	0.07 (0.67)
2D	0.09 (0.96)	0.03 (0.99)	<0.01 (0.73)	0.02 (0.99)	0.20 (0.98)
norm. $p_T^{\nu\nu}$	0.60 (0.87)	0.40 (0.92)	0.60 (0.67)	0.57 (0.98)	0.10 (0.70)
norm. $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$	0.21 (0.22)	0.20 (0.24)	0.10 (0.14)	0.04 (0.37)	0.04 (0.33)
norm. 2D	0.08 (0.14)	0.03 (0.21)	<0.01 (0.02)	0.02 (0.80)	0.19 (0.67)

of most predictions to below unity. Only the MC@NLO+PYTHIA prediction leads to χ^2/ndf values significantly above unity for the normalised 2D measurement. The results of the χ^2 tests without including uncertainties show that POWHEG+PYTHIA yields overall the best description for the two one-dimensional absolute differential measurements, and the NNLO calculation shows the best agreement with the 2D absolute result.

For the normalized differential cross sections, shown in Fig. 8, the systematic uncertainties are slightly smaller than those for the absolute cross section measurements. This especially holds for the $\min[\Delta\phi(\vec{p}_T^{\nu\nu}, \vec{p}_T^\ell)]$ measurement, where the dominant uncertainty on the absolute differential measurements that arises from lepton reconstruction efficiency is significantly reduced, which emphasizes the shape differences between the measurement and predictions already discussed for the absolute result. Similar to the absolute results, all predictions are in reasonable agreement with the measurements as shown by the obtained χ^2 values.

8 Summary

Differential cross section measurements have been presented for the top quark pair production in the dileptonic channel in proton-proton (pp) collisions at a center-of-mass energy of 13 TeV using observables based on the dineutrino kinematic properties. The measurements are

performed based on the pp collision data recorded by the CMS detector at the CERN LHC between 2016 and 2018, corresponding to an integrated luminosity of 138 fb^{-1} .

The differential cross sections are measured as functions of the transverse momentum of the dineutrino system p_T^{vv} , and the minimal azimuthal angle between the dineutrino system and a charged lepton $\min[\Delta\phi(\vec{p}_T^{vv}, \vec{p}_T^\ell)]$, as well as both observables in two dimensions. To improve the resolution of the missing transverse momentum, which serves as a measure of p_T^{vv} in signal events, a dedicated deep neural network regression has been developed. The method significantly improves the resolution of both the magnitude and the azimuthal angle of the missing transverse momentum. The absolute and normalized differential cross section results are obtained based on an unregularized least squares unfolding method.

The differential cross sections are compared to predictions based on Monte Carlo simulation as well as two fixed-order theoretical calculations, corresponding to next-to-leading order and next-to-next-to-leading order (NNLO) accuracy in quantum chromodynamics. A remarkable agreement between the different theoretical predictions and the measured differential cross sections is observed. For both one-dimensional measurements, the best overall description is provided by the two POWHEG predictions, while for the two-dimensional measurement the best agreement is observed for the NNLO fixed-order calculation. However, the differences between the five predictions are mostly small, such that none of the predictions is significantly disfavored by the measured differential cross sections. These results constitute the first differential cross section measurements based on the dineutrino kinematic properties in top quark pair production.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leven-

tis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA) and Fonds voor Wetenschappelijk Onderzoek contract No. 1228724N (Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, and the Natural Science Foundation of China under Grant No. 12061141002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ.KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR – Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, the National Science, Research and Innovation Fund program IND_FF_68.369.2300.097, and the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] ATLAS Collaboration, “Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **73** (2013) 2261, doi:10.1140/epjc/s10052-012-2261-1, arXiv:1207.5644.
- [2] CMS Collaboration, “Measurement of differential top-quark pair production cross sections in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **73** (2013) 2339, doi:10.1140/epjc/s10052-013-2339-4, arXiv:1211.2220.
- [3] ATLAS Collaboration, “Measurements of normalized differential cross sections for $t\bar{t}$ production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, *Phys. Rev. D* **90** (2014) 072004, doi:10.1103/PhysRevD.90.072004, arXiv:1407.0371.
- [4] ATLAS Collaboration, “Differential top-antitop cross-section measurements as a function of observables constructed from final-state particles using pp collisions at $\sqrt{s} = 7$ TeV in the ATLAS detector”, *JHEP* **06** (2015) 100, doi:10.1007/JHEP06(2015)100, arXiv:1502.05923.

-
- [5] ATLAS Collaboration, “Measurement of top quark pair differential cross-sections in the dilepton channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS”, *Phys. Rev. D* **94** (2016) 092003, doi:10.1103/PhysRevD.94.092003, arXiv:1607.07281.
[Addendum: doi:10.1103/PhysRevD.101.119901].
- [6] CMS Collaboration, “Measurement of the differential cross section for top quark pair production in pp collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **75** (2015) 542, doi:10.1140/epjc/s10052-015-3709-x, arXiv:1505.04480.
- [7] ATLAS Collaboration, “Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **76** (2016) 538, doi:10.1140/epjc/s10052-016-4366-4, arXiv:1511.04716.
- [8] ATLAS Collaboration, “Measurement of the differential cross-section of highly boosted top quarks as a function of their transverse momentum in $\sqrt{s} = 8$ TeV proton-proton collisions using the ATLAS detector”, *Phys. Rev. D* **93** (2016) 032009, doi:10.1103/PhysRevD.93.032009, arXiv:1510.03818.
- [9] CMS Collaboration, “Measurement of the $t\bar{t}$ production cross section in the all-jets final state in pp collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76** (2016) 128, doi:10.1140/epjc/s10052-016-3956-5, arXiv:1509.06076.
- [10] CMS Collaboration, “Measurement of the integrated and differential $t\bar{t}$ production cross sections for high- p_T top quarks in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **94** (2016) 072002, doi:10.1103/PhysRevD.94.072002, arXiv:1605.00116.
- [11] CMS Collaboration, “Measurement of double-differential cross sections for top quark pair production in pp collisions at $\sqrt{s} = 8$ TeV and impact on parton distribution functions”, *Eur. Phys. J. C* **77** (2017) 459, doi:10.1140/epjc/s10052-017-4984-5, arXiv:1703.01630.
- [12] ATLAS Collaboration, “Measurement of the top-quark mass in $t\bar{t}+1$ -jet events collected with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **11** (2019) 150, doi:10.1007/JHEP11(2019)150, arXiv:1905.02302.
- [13] CMS Collaboration, “Measurement of differential cross sections for top quark pair production using the lepton+jets final state in proton-proton collisions at 13 TeV”, *Phys. Rev. D* **95** (2017) 092001, doi:10.1103/PhysRevD.95.092001, arXiv:1610.04191.
- [14] ATLAS Collaboration, “Measurement of jet activity produced in top-quark events with an electron, a muon and two b-tagged jets in the final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **77** (2017) 220, doi:10.1140/epjc/s10052-017-4766-0, arXiv:1610.09978.
- [15] ATLAS Collaboration, “Measurements of top-quark pair differential cross-sections in the $e\mu$ channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **77** (2017) 292, doi:10.1140/epjc/s10052-017-4821-x, arXiv:1612.05220.
- [16] CMS Collaboration, “Measurement of normalized differential $t\bar{t}$ cross sections in the dilepton channel from pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **04** (2018) 060, doi:10.1007/JHEP04(2018)060, arXiv:1708.07638.

- [17] CMS Collaboration, “Measurement of differential cross sections for the production of top quark pairs and of additional jets in lepton+jets events from pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **97** (2018) 112003, doi:10.1103/PhysRevD.97.112003, arXiv:1803.08856.
- [18] CMS Collaboration, “Measurements of $t\bar{t}$ differential cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV using events containing two leptons”, *JHEP* **02** (2019) 149, doi:10.1007/JHEP02(2019)149, arXiv:1811.06625.
- [19] CMS Collaboration, “Measurement of $t\bar{t}$ normalised multi-differential cross sections in pp collisions at $\sqrt{s} = 13$ TeV, and simultaneous determination of the strong coupling strength, top quark pole mass, and parton distribution functions”, *Eur. Phys. J. C* **80** (2020) 658, doi:10.1140/epjc/s10052-020-7917-7, arXiv:1904.05237.
- [20] ATLAS Collaboration, “Measurements of top-quark pair differential and double-differential cross-sections in the ℓ +jets channel with pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Eur. Phys. J. C* **79** (2019) 1028, doi:10.1140/epjc/s10052-019-7525-6, arXiv:1908.07305. [Erratum: doi:10.1140/epjc/s10052-020-08541-3].
- [21] ATLAS Collaboration, “Measurement of the $t\bar{t}$ production cross-section and lepton differential distributions in $e\mu$ dilepton events from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **80** (2020) 528, doi:10.1140/epjc/s10052-020-7907-9, arXiv:1910.08819.
- [22] ATLAS Collaboration, “Measurements of top-quark pair single- and double-differential cross-sections in the all-hadronic channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *JHEP* **01** (2021) 033, doi:10.1007/JHEP01(2021)033, arXiv:2006.09274.
- [23] CMS Collaboration, “Measurement of differential $t\bar{t}$ production cross sections in the full kinematic range using lepton+jets events from proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **104** (2021) 092013, doi:10.1103/PhysRevD.104.092013, arXiv:2108.02803.
- [24] S. Weinberg, “Supersymmetry at ordinary energies. Masses and conservation laws”, *Phys. Rev. D* **26** (1982) 287, doi:10.1103/PhysRevD.26.287.
- [25] W. Beenakker et al., “NNLL resummation for stop pair-production at the LHC”, *JHEP* **05** (2016) 153, doi:10.1007/JHEP05(2016)153, arXiv:1601.02954.
- [26] CMS Collaboration, “Search for top squark pair production using dilepton final states in pp collision data collected at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **81** (2021) 3, doi:10.1140/epjc/s10052-020-08701-5, arXiv:2008.05936.
- [27] CMS Collaboration, “Combined searches for the production of supersymmetric top quark partners in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **81** (2021) 970, doi:10.1140/epjc/s10052-021-09721-5, arXiv:2107.10892.
- [28] M. Czakon, A. Mitov, and R. Poncelet, “NNLO QCD corrections to leptonic observables in top-quark pair production and decay”, *JHEP* **05** (2021) 212, doi:10.1007/JHEP05(2021)212, arXiv:2008.11133.
- [29] HEPData record for this analysis, 2025. doi:10.17182/hepdata.153302.

-
- [30] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [31] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [32] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [33] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [34] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [35] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [36] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [37] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [38] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [39] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [40] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [41] NNPDF Collaboration, “Unbiased global determination of parton distributions and their uncertainties at NNLO and at LO”, *Nucl. Phys. B* **855** (2012) 153, doi:10.1016/j.nuclphysb.2011.09.024, arXiv:1107.2652.
- [42] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
- [43] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [44] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA 8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.

- [45] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [46] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [47] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [48] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [49] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [50] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [51] J. Bellm et al., “HERWIG 7.0/HERWIG++ 3.0 release note”, *Eur. Phys. J. C* **76** (2016) 196, doi:10.1140/epjc/s10052-016-4018-8, arXiv:1512.01178.
- [52] CMS Collaboration, “Development and validation of HERWIG 7 tunes from CMS underlying-event measurements”, *Eur. Phys. J. C* **81** (2021) 312, doi:10.1140/epjc/s10052-021-08949-5, arXiv:2011.03422.
- [53] J. Alwall et al., “Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions”, *Eur. Phys. J. C* **53** (2008) 473, doi:10.1140/epjc/s10052-007-0490-5, arXiv:0706.2569.
- [54] S. Alioli, P. Nason, C. Oleari, and E. Re, “NLO single-top production matched with shower in POWHEG: s - and t -channel contributions”, *JHEP* **09** (2009) 111, doi:10.1088/1126-6708/2009/09/111, arXiv:0907.4076. [Erratum: doi:10.1007/JHEP02(2010)011].
- [55] E. Re, “Single-top Wt -channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [56] M. Czakon and A. Mitov, “TOP++: a program for the calculation of the top-pair cross-section at hadron colliders”, *Comput. Phys. Commun.* **185** (2014) 2930, doi:10.1016/j.cpc.2014.06.021, arXiv:1112.5675.
- [57] M. Czakon, P. Fiedler, and A. Mitov, “Total top-quark pair-production cross section at hadron colliders through $\mathcal{O}(\alpha_s^4)$ ”, *Phys. Rev. Lett.* **110** (2013) 252004, doi:10.1103/PhysRevLett.110.252004, arXiv:1303.6254.
- [58] M. Cacciari et al., “Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation”, *Phys. Lett. B* **710** (2012) 612, doi:10.1016/j.physletb.2012.03.013, arXiv:1111.5869.

-
- [59] P. Bärnreuther, M. Czakon, and A. Mitov, “Percent level precision physics at the Tevatron: Next-to-next-to-leading order QCD corrections to $q\bar{q} \rightarrow t\bar{t} + X$ ”, *Phys. Rev. Lett.* **109** (2012) 132001, doi:10.1103/PhysRevLett.109.132001, arXiv:1204.5201.
 - [60] M. Czakon and A. Mitov, “NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels”, *JHEP* **12** (2012) 054, doi:10.1007/JHEP12(2012)054, arXiv:1207.0236.
 - [61] M. Czakon et al., “Top-pair production at the LHC through NNLO QCD and NLO EW”, *JHEP* **10** (2017) 186, doi:10.1007/JHEP10(2017)186, arXiv:1705.04105.
 - [62] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “FEWZ 2.0: A code for hadronic Z production at next-to-next-to-leading order”, *Comput. Phys. Commun.* **182** (2011) 2388, doi:10.1016/j.cpc.2011.06.008, arXiv:1011.3540.
 - [63] Y. Li and F. Petriello, “Combining QCD and electroweak corrections to dilepton production in FEWZ”, *Phys. Rev. D* **86** (2012) 094034, doi:10.1103/PhysRevD.86.094034, arXiv:1208.5967.
 - [64] N. Kidonakis, “Two-loop soft anomalous dimensions for single top quark associated production with a W^- or H^- ”, *Phys. Rev. D* **82** (2010) 054018, doi:10.1103/PhysRevD.82.054018, arXiv:1005.4451.
 - [65] M. Aliev et al., “HATHOR: Hadronic top and heavy quarks cross section calculator”, *Comput. Phys. Commun.* **182** (2011) 1034, doi:10.1016/j.cpc.2010.12.040, arXiv:1007.1327.
 - [66] P. Kant et al., “HATHOR for single top-quark production: Updated predictions and uncertainty estimates for single top-quark production in hadronic collisions”, *Comput. Phys. Commun.* **191** (2015) 74, doi:10.1016/j.cpc.2015.02.001, arXiv:1406.4403.
 - [67] T. Gehrmann et al., “ W^+W^- production at hadron colliders in next to next to leading order QCD”, *Phys. Rev. Lett.* **113** (2014) 212001, doi:10.1103/PhysRevLett.113.212001, arXiv:1408.5243.
 - [68] J. M. Campbell, R. K. Ellis, and C. Williams, “Vector boson pair production at the LHC”, *JHEP* **07** (2011) 018, doi:10.1007/JHEP07(2011)018, arXiv:1105.0020.
 - [69] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015. doi:10.17181/CERN.VU8I.D59J.
 - [70] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
 - [71] CMS Collaboration, “Performance of the CMS electromagnetic calorimeter in pp collisions at $\sqrt{s} = 13$ TeV”, *JINST* **19** (2024) P09004, doi:10.1088/1748-0221/19/09/P09004, arXiv:2403.15518.
 - [72] CMS Collaboration, “Performance of CMS muon reconstruction from proton-proton to heavy ion collisions”, *JINST* **19** (2024) P09012, doi:10.1088/1748-0221/19/09/P09012, arXiv:2404.17377.

- [73] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [74] M. Cacciari, G. P. Salam, and G. Soyez, “FASTJET user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [75] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [76] E. Bols et al., “Jet flavour classification using DeepJet”, *JINST* **15** (2020) P12012, doi:10.1088/1748-0221/15/12/P12012, arXiv:2008.10519.
- [77] CMS Collaboration, “Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector”, *JINST* **14** (2019) P07004, doi:10.1088/1748-0221/14/07/P07004, arXiv:1903.06078.
- [78] D. Bertolini, P. Harris, M. Low, and N. Tran, “Pileup per particle identification”, *JHEP* **10** (2014) 059, doi:10.1007/JHEP10(2014)059, arXiv:1407.6013.
- [79] Particle Data Group, S. Navas et al., “Review of particle physics”, *Phys. Rev. D* **110** (2024) 030001, doi:10.1103/PhysRevD.110.030001.
- [80] CMS Collaboration, “Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV”, *JINST* **13** (2018) P05011, doi:10.1088/1748-0221/13/05/P05011, arXiv:1712.07158.
- [81] M. Abadi et al., “TENSORFLOW: Large-scale machine learning on heterogeneous systems”, 2015. Software available from <http://tensorflow.org>. doi:10.5281/zenodo.4724125.
- [82] F. Chollet et al., “KERAS”, 2015. Software available from <https://keras.io>.
- [83] R. Cousins, “Generalization of chisquare goodness-of-fit test for binned data using saturated models, with application to histograms”, 2013.
- [84] CMS Collaboration, “The CMS statistical analysis and combination tool: COMBINE”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [85] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [86] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [87] CMS Collaboration, “Differential cross section measurements for the production of top quark pairs and of additional jets using dilepton events from pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **02** (2025) 064, doi:10.1007/JHEP02(2025)064, arXiv:2402.08486.
- [88] A. Andreassen and B. Nachman, “Neural networks for full phase-space reweighting and parameter tuning”, *Phys. Rev. D* **101** (2020) 091901, doi:10.1103/PhysRevD.101.091901, arXiv:1907.08209.

- [89] CMS Collaboration, “Reweighting simulated events using machine-learning techniques in the CMS experiment”, *Eur. Phys. J. C* **85** (2025) 495, doi:10.1140/epjc/s10052-025-14097-x, arXiv:2411.03023.
- [90] CMS Collaboration, “CMS PYTHIA 8 colour reconnection tunes based on underlying-event data”, *Eur. Phys. J. C* **83** (2023) 587, doi:10.1140/epjc/s10052-023-11630-8, arXiv:2205.02905.
- [91] S. Schmitt, “TUNFOLD, an algorithm for correcting migration effects in high energy physics”, *JINST* **7** (2012) T10003, doi:10.1088/1748-0221/7/10/T10003, arXiv:1205.6201.
- [92] ATLAS Collaboration, “Inclusive and differential cross-sections for dilepton $t\bar{t}$ production measured in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *JHEP* **07** (2023) 141, doi:10.1007/JHEP07(2023)141, arXiv:2303.15340.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

V. Chekhovsky, A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck² , R. Schöffbeck² , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz² 

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , L. Lambrecht , D. Marckx , K. Skovpen , N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote⁵ , A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶ , E.M. Da Costa , G.G. Da Silveira⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza , T. Laux Kuhn⁷ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu

Institute of High Energy Physics, Beijing, China

G.M. Chen⁸ , H.S. Chen⁸ , M. Chen⁸ , Q. Hou , F. Iemmi , C.H. Jiang, A. Kapoor⁹ ,
H. Liao , Z.-A. Liu¹⁰ , R. Sharma¹¹ , J.N. Song¹⁰, J. Tao , C. Wang⁸, J. Wang ,
H. Zhang , J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang ,
A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang,
Y. Zhao, C. Zhou 

State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹², B. Li¹³, H. Wang , K. Yi¹⁴ , J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

J. Jaramillo , C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov ,
T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos , F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A.A. Abdelalim^{15,16} , S. Elgammal¹⁷, A. Ellithi Kamel¹⁸ 

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad , M.A. Mahmoud 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki , M.m. Rantanen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

H. Kirschenmann , P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure , F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²⁰ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules,

CNRS/IN2P3, Villeurbanne, FranceA. Di Florio **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**

D. Amram , S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,
 P. Depasse , C. Dozen²¹ , H. El Mamouni , J. Fay , S. Gascon , M. Gouzevitch ,
 C. Greenberg , G. Grenier , B. Ille , E. Jourdhuy , I.B. Laktineh , M. Lethuillier ,
 L. Mirabito , S. Perries , A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, GeorgiaI. Lomidze , T. Toriashvili²² , Z. Tsamalaidze²³ **RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,
 P. Nattland , A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann ,
 M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung ,
 N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , F. Nowotny ,
 A. Pozdnyakov , Y. Rath , W. Redjeb , F. Rehm , H. Reithler , V. Sarkisovi , A. Schmidt ,
 C. Seth , A. Sharma , J.L. Spah , F. Torres Da Silva De Araujo²⁴ , S. Wiedenbeck ,
 S. Zaleski 

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons ,
 A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso , Y. An , J. Bach ,
 S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere ,
 F. Blekman²⁵ , K. Borras²⁶ , A. Campbell , A. Cardini , S. Chatterjee , F. Colombina ,
 M. De Silva , G. Eckerlin , D. Eckstein , E. Gallo²⁵ , A. Geiser , V. Guglielmi ,
 M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort ,
 R. Kogler , M. Komm , D. Krücker , W. Lange , D. Leyva Pernia , K. Lipka²⁷ ,
 W. Lohmann²⁸ , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin ,
 A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller ,
 L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza ,
 D. Rastorguev , J. Rübenach , L. Rygaard , M. Scham^{29,26} , S. Schnake²⁶ , P. Schütze ,
 C. Schwanenberger²⁵ , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford ,
 F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann ,
 L. Wiens²⁶ , C. Wissing , Y. Yang , S. Zakharov , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bollweg , M. Bonanomi , P. Connor ,
 K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen ,
 H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer ,
 C.c. Kuo , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux ,
 M. Mrowietz , A. Nigamova , K. Nikolopoulos , Y. Nissan , A. Paasch , K.J. Pena Rodriguez ,
 T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper ,
 M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews ,
 B. Wiederspan , M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann³⁰ , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa , L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou , G. Daskalakis , A. Kyriakis , A. Papadopoulos³⁰ , A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos

National Technical University of Athens, Athens, Greece

T. Chatzistavrou , G. Karapostoli , K. Kousouris , E. Siamarkou, G. Tsiopolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath^{31,32} , K. Márton, A.J. Rádl³³ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³⁴ , M.M.A. Gadallah³⁵ , Á. Kadlecik , G. Pásztor , G.I. Veres

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³⁴ , F. Nemes³⁴ , T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁶ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla

University of Delhi, Delhi, India

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya

Indian Institute of Technology Kanpur, Kanpur, India

S. Mukherjee 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman³⁷ , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁸ , D. Kumar³⁸ , T. Mishra , B. Parida³⁹ , P.R. Pujahari , N.R. Saha , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad , G.B. Mohanty , M. Shelake , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , S. Bhowmik⁴⁰ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati⁴¹ , C. Kar , D. Maity⁴² , P. Mal , K. Naskar⁴² , A. Nayak⁴² , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese⁴² , D. Vats⁴² 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya⁴³ , A. Alpana , S. Dube , B. Gomber⁴³ , P. Hazarika , B. Kansal , A. Laha , B. Sahu⁴³ , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴⁴ , A. Jafari⁴⁵ , M. Zeinali⁴⁶ 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri , S. Chenarani⁴⁷ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁸ 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , B. D'Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,15} , N. Ferrara^{a,b} , L. Fiore^a , G. Iaselli^{a,c} , L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellicchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} 

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , A. Castro^{+a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,49} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,50} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,50} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^a , G. Barbagli^a , G. Bardelli^{a,b} , M. Bartolini^{a,b} , B. Camaiani^{a,b} 

A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , L. Damenti^{a,b} , E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁵¹ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , G. Lavizzari^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , B.S. Pinolini^a , G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b} , N. Cavallo^{a,c} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,52} , P. Paolucci^{a,30} , B. Rossi^a 

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,53} , D. Bisello^{a,b} , P. Bortignon^a , G. Bortolato^{a,b} , A.C.M. Bulla^a , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^{a,54} , F. Fanzago^a , U. Gasparini^{a,b} , S. Giorgetti^a , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , D. Fiorina^a , P. Montagna^{a,b} , M. Pelliccioni^a , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,55} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^{a,b} , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^{a,b} , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b,42} , M. Musich^{a,b} , S. Nandan^a , F. Palla^a , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,40} , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} 

E. Di Marco^{a,b} , M. Diemoz^a , F. Errico^{a,b} , R. Gargiulo^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , P. De Remigis^a , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , L. Markovic^{a,b} , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, D. Lee, H. Lee , S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNU, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi⁵⁶ , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou 

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , S. Yang 

Sejong University, Seoul, Korea

Y. Kang , H. S. Kim , Y. Kim , S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , K. Hwang , B. Kim , K. Lee , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM),

Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs , A. Potrebko ,
M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁷ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma , E. De La Cruz-Burelo , I. Heredia-
De La Cruz⁵⁸ , R. Lopez-Fernandez , J. Mejia Guisao , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar , A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 

AGH University of Krakow, Krakow, Poland

V. Avati, A. Bellora⁵⁹ , L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,
A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo ,
T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo 

G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic, M. Stojanovic 

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Sastre , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , E. Palencia Cortezon , J. Prado Pico , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasoa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁶⁰ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁶¹ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo , E. Brondolin , C.E. Brown , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, J. Hegeman , J.K. Heikkilä , B. Huber , V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁸ , G. Karathanasis , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , M. Magherini , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , M. Migliorini , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabadý , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁶² , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter 

J. Wanczyk⁶³ , J. Wang, S. Wuchterl , P. Zehetner , P. Zejdl , W.D. Zeuner 

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶⁴ , L. Caminada⁶⁴ , A. Ebrahimi , W. Erdmann , R. Horisberger ,
Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶⁴ , L. Noehte⁶⁴ ,
T. Rohe , A. Samalan 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga ,
K. Datta , P. De Bryas Dexmiers D'archiac⁶³ , A. De Cosa , G. Dissertori , M. Dittmar,
M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer ,
T.G. Harte , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani ,
L. Marchese , A. Mascellani⁶³ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini ,
B. Ristic , R. Seidita , J. Steggemann⁶³ , A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler⁶⁵ , P. Bäertschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin ,
A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liehti , A. Macchiolo , P. Meiring ,
F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr , F. Stäger ,
R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶⁶, D. Bhowmik, C.M. Kuo, W. Lin , P.K. Rout , P.C. Tiwari³⁸ 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao,
S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker ,
L.s. Tsai, D. Tsiou, H.y. Wu , E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu⁶⁷ , E. Eskut , Y. Guler⁶⁸ ,
E. Gulpinar Guler⁶⁸ , C. Isik , O. Kara , A. Kayis Topaksu , Y. Komurcu ,
G. Onengut , K. Ozdemir⁶⁹ , A. Polatoz , B. Tali⁷⁰ , U.G. Tok , E. Uslan ,
I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁷¹ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁷² , O. Kaya⁷³ , S. Tekten⁷⁴ 

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{67,75} , S. Sen⁷⁶ 

Istanbul University, Istanbul, Turkey

O. Aydilek⁷⁷ , B. Haciosahinoglu , I. Hos⁷⁸ , B. Kaynak , S. Ozkorucuklu , O. Potok ,
H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci , B. Isildak⁷⁹ , D. Sunar Cerci , T. Yetkin 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, V.J. Smith , K. Walkingshaw Pass

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁸⁰ , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya , G.S. Chahal⁸¹ , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , A. Howard, G. Iles , C.R. Knight , P. Krueper , J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios, M. Mieskolainen , J. Nash⁸² , M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards, A. Rose , L. Russell , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee³⁰ , M. Vojinovic , N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu 

The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio⁸³ , E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA

G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , J.M. Hogan⁸⁴ , T. Kwon , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , T. Russell , S. Sagir⁸⁵ , X. Shen , M. Stamenkovic , N. Venkatasubramanian 

University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , S. Yoo 

University of California, Los Angeles, California, USA

K. Adamidis , M. Bachtis , D. Campos , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca , A. Nunez Del Prado , D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch , M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , M.M. Hussain , J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang 

California Institute of Technology, Pasadena, California, USA

A. Albert , S. Bhattacharya , A. Bornheim , O. Cerri , J. Mao , H.B. Newman , G. Reales Gutiérrez , M. Spiropulu , J.R. Vlimant , S. Xie , R.Y. Zhu 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi , V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil , M. Kanemura , C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , N. Manganello , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , D.J. Cranshaw , J. Dickinson , J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives , J. Monroy , G. Niendorf , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green , A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma 

B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸⁶ , F. Ravera , A. Reinsvold Hall⁸⁷ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field , C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez , H. Prosper , P.R. Prova , M. Wulansatiti , R. Yohay , J. Zhang 

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , T. Elkafrawy⁸⁸ , M. Hohlmann , E. Yanes 

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , A. Baty , C. Bennett , R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth , A. Hingrajiya , D.J. Hofman , J.h. Lee , D. S. Lemos , C. Mills , S. Nanda , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti , P. Roy , T. Roy , S. Rudrabhatla , N. Singh , M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend , K. Dilsiz⁸⁹ , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo , A. Mestvirishvili⁹⁰ , O. Neogi , H. Ogul⁹¹ , Y. Onel , A. Penzo , C. Snyder , E. Tiras⁹² 

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁹³ , C. Rogan , C. Royon , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim , S. Lascio , P. Major , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe , E. Popova⁹⁴ , A. Shevelev , L. Wang , L. Zhang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , J. Bendavid , S. Bright-Thonney , I.A. Cali , P.c. Chou 

M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso , P. Harris, D. Hoang , D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , P. Levchenko , J. Li , M. Lu , R. McCarthy , D.M. Morse , T. Orimoto , A. Parker , L. Skinnari , E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , D. Li , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco 

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²³ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalá 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, D.A. Wenzl, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , P. Fackeldey , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , Y. Lai , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully , L.H. Vage 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma 

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , J.F. Schulte , A. K. Viridi , F. Wang , A. Wildridge , W. Xie , Y. Yao 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar , P. Parygin⁹⁴ , R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath , J. Martins , R. Montalvo, K. Nash, J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov⁶³ , A. Bolshov, O. Bouhali⁹⁵ , R. Eusebi , J. Gilmore , T. Huang , T. Kamon⁹⁶ , H. Kim , S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane , S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez 

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna , A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , M. Danilov⁹⁷ , T. Dimova⁹⁷ , A. Ershov⁹⁷ , I. Golutvin[†] , I. Gorbunov , A. Gribushin⁹⁷ , V. Karjavine , V. Klyukhin⁹⁷ , O. Kodolova^{98,94} , V. Korenkov , A. Kozyrev⁹⁷ , A. Lanev , A. Malakhov , V. Matveev⁹⁷ , A. Nikitenko^{99,98} , V. Palichik , V. Perelygin , S. Petrushanko⁹⁷ , O. Radchenko⁹⁷ , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹⁷ , V. Smirnov , O. Teryaev , I. Tlisova⁹⁷ , A. Toropin , N. Voytishin , B.S. Yuldashev^{†100}, A. Zarubin , I. Zhizhin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

G. Gavrillov , V. Golovtsov , Y. Ivanov , V. Kim⁹⁷ , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†], A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , K. Ivanov , V. Gavrillov , N. Lychkovskaya , V. Popov , A. Zhokin , R. Chistov⁹⁷ , S. Polikarpov⁹⁷ , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , V. Bunichev , L. Dudko , M. Perfilov , V. Savrin , G. Vorotnikov , V. Blinov⁹⁷, V. Kachanov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at University of Chinese Academy of Sciences, Beijing, China

⁹Also at China Center of Advanced Science and Technology, Beijing, China

¹⁰Also at University of Chinese Academy of Sciences, Beijing, China

¹¹Also at China Spallation Neutron Source, Guangdong, China

¹²Now at Henan Normal University, Xinxiang, China

¹³Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁴Now at The University of Iowa, Iowa City, Iowa, USA

¹⁵Also at Helwan University, Cairo, Egypt

¹⁶Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁷Now at British University in Egypt, Cairo, Egypt

¹⁸Now at Cairo University, Cairo, Egypt

¹⁹Also at Purdue University, West Lafayette, Indiana, USA

²⁰Also at Université de Haute Alsace, Mulhouse, France

²¹Also at Istinye University, Istanbul, Turkey

²²Also at Tbilisi State University, Tbilisi, Georgia

²³Also at an institute formerly covered by a cooperation agreement with CERN

²⁴Also at The University of the State of Amazonas, Manaus, Brazil

²⁵Also at University of Hamburg, Hamburg, Germany

²⁶Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁷Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁸Also at Brandenburg University of Technology, Cottbus, Germany

²⁹Also at Forschungszentrum Jülich, Juelich, Germany

³⁰Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³¹Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³²Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³³Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³⁴Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³⁵Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁶Also at Punjab Agricultural University, Ludhiana, India

³⁷Also at University of Visva-Bharati, Santiniketan, India

³⁸Also at Indian Institute of Science (IISc), Bangalore, India

³⁹Also at Amity University Uttar Pradesh, Noida, India

- ⁴⁰ Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁴¹ Also at IIT Bhubaneswar, Bhubaneswar, India
- ⁴² Also at Institute of Physics, Bhubaneswar, India
- ⁴³ Also at University of Hyderabad, Hyderabad, India
- ⁴⁴ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴⁵ Also at Isfahan University of Technology, Isfahan, Iran
- ⁴⁶ Also at Sharif University of Technology, Tehran, Iran
- ⁴⁷ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁸ Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁹ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁵⁰ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁵¹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵² Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵³ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵⁴ Also at Lulea University of Technology, Lulea, Sweden
- ⁵⁵ Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁶ Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁷ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁸ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁹ Also at INFN Sezione di Torino, Università di Torino, Torino, Italy; Università del Piemonte Orientale, Novara, Italy
- ⁶⁰ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁶¹ Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁶² Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶³ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶⁴ Also at Universität Zürich, Zurich, Switzerland
- ⁶⁵ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁶ Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁷ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁸ Also at Konya Technical University, Konya, Turkey
- ⁶⁹ Also at Izmir Bakircay University, Izmir, Turkey
- ⁷⁰ Also at Adiyaman University, Adiyaman, Turkey
- ⁷¹ Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁷² Also at Marmara University, Istanbul, Turkey
- ⁷³ Also at Milli Savunma University, Istanbul, Turkey
- ⁷⁴ Also at Kafkas University, Kars, Turkey
- ⁷⁵ Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁶ Also at Hacettepe University, Ankara, Turkey
- ⁷⁷ Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁸ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁹ Also at Yildiz Technical University, Istanbul, Turkey
- ⁸⁰ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁸¹ Also at IPPP Durham University, Durham, United Kingdom

⁸²Also at Monash University, Faculty of Science, Clayton, Australia

⁸³Also at Università di Torino, Torino, Italy

⁸⁴Also at Bethel University, St. Paul, Minnesota, USA

⁸⁵Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁸⁶Also at California Institute of Technology, Pasadena, California, USA

⁸⁷Also at United States Naval Academy, Annapolis, Maryland, USA

⁸⁸Also at Ain Shams University, Cairo, Egypt

⁸⁹Also at Bingol University, Bingol, Turkey

⁹⁰Also at Georgian Technical University, Tbilisi, Georgia

⁹¹Also at Sinop University, Sinop, Turkey

⁹²Also at Erciyes University, Kayseri, Turkey

⁹³Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹⁴Now at another institute formerly covered by a cooperation agreement with CERN

⁹⁵Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁶Also at Kyungpook National University, Daegu, Korea

⁹⁷Also at another institute formerly covered by a cooperation agreement with CERN

⁹⁸Also at Yerevan Physics Institute, Yerevan, Armenia

⁹⁹Also at Imperial College, London, United Kingdom

¹⁰⁰Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan