CERN-EP-2025-205
2025/10/23

CMS-TOP-24-009

Observation of tWZ production at the CMS experiment

The CMS Collaboration*

Abstract

The first observation of single top quark production in association with a W and a Z boson in proton-proton collisions is reported. The analysis uses data at center-of-mass energies of 13 and 13.6 TeV recorded with the CMS detector at the CERN LHC, corresponding to a total integrated luminosity of 200 fb^{-1} . Events with three or four charged leptons, which can be electrons or muons, are selected. Advanced machine-learning algorithms and improved reconstruction methods, compared to an earlier analysis, result in an unprecedented sensitivity to tWZ production. The measured cross sections for tWZ production are $248 \pm 52 \text{ fb}$ and $244 \pm 74 \text{ fb}$ for $\sqrt{s} = 13$ and 13.6 TeV, respectively. The signal is established with a statistical significance of 5.8 standard deviations, with 3.5 expected, compared to the background-only hypothesis.

Submitted to Physical Review Letters

The standard model (SM) of particle physics provides a remarkably successful theoretical framework for describing the fundamental particles and their interactions through the electromagnetic, weak, and strong forces. Over the past several decades, the predictions of the SM have been extensively tested with high precision, and its theoretical consistency has withstood rigorous experimental scrutiny. Despite its successes, the SM is known to be an incomplete theory. For example, it does not account for dark matter, the observed matter-antimatter asymmetry in the Universe, neutrino masses, and the quantum nature of gravity. Consequently, the search for phenomena not included in the SM remains a central focus of contemporary particle physics.

High sensitivity to deviations from the SM predictions, and therefore to beyond-the-SM (BSM) effects, can be achieved in direct searches [1, 2], precision measurements [3], studies of loop-induced processes [4], or through measurements of very rare processes. Among the rare processes predicted by the SM, the production of a single top quark t in association with a W and a Z boson in proton-proton (pp) collisions, $pp \rightarrow tWZ$, is particularly interesting. This process is exceptionally rare with predicted cross sections of 136^{+9}_{-8} fb at $\sqrt{s} = 13$ TeV and 148^{+10}_{-9} fb at 13.6 TeV at next-to-leading order (NLO) accuracy in quantum chromodynamics (QCD) [5, 6]. The tWZ process provides direct access to the electroweak (EW) couplings of the top quark through Feynman diagrams similar to the example shown in Fig. 1 (left). The tWZ production allows for stringent tests of the gauge structure of the SM and the underlying symmetries governing particle interactions [7, 8]. New heavy particles can induce anomalous top-quark couplings with the EW gauge bosons, as shown by the Feynman diagram in Fig. 1 (right). These would manifest as measurable deviations in the rate of tWZ production, making measurements thereof a sensitive probe of BSM physics [9, 10].

This Letter presents an analysis of pp collision data from the CERN LHC, aiming to identify events from tWZ production, characterized by the presence of multiple leptons, jets, and missing transverse momentum (p_T^{miss}). However, the experimental challenges are significant. The small production cross section of the tWZ process compared to backgrounds with similar signature, namely $t\bar{t}Z$ and diboson production with additional jets, necessitates advanced analysis techniques to isolate the signal. Evidence for tWZ production has been reported by the CMS Collaboration in 2024 using $\sqrt{s} = 13$ TeV data corresponding to an integrated luminosity of 138 fb^{-1} , recorded in 2016–2018 [11]. The previous analysis reported an observed statistical significance of 3.4 standard deviations (s.d.) with 1.4 s.d. expected. The analysis presented in this Letter uses state-of-the-art machine-learning (ML) algorithms and improved reconstruction methods to study tWZ production with unprecedented sensitivity. It updates the analysis of 13 TeV data and includes, in addition, pp collision data at 13.6 TeV recorded in 2022–2023. The total integrated luminosity of the dataset is 200 fb^{-1} , of which 62 fb^{-1} were recorded at 13.6 TeV. Tabulated results are provided in the HEPData record for this analysis [12].




Figure 1: Example Feynman diagrams for tWZ production in the SM (left) and for anomalous tWZ production through an effective interaction (right). Charge conjugate versions of these diagrams are implied.

The CMS apparatus [13, 14] is a multipurpose, nearly hermetic detector, designed to trigger on [15–17] and identify electrons, muons, photons, and hadrons [18–20]. A global “particle-flow” (PF) algorithm [21] aims to reconstruct all individual particles in an event, combining information provided by the all-silicon inner tracker and by the crystal electromagnetic (ECAL) and brass-scintillator hadron calorimeters (HCAL), operating inside a 3.8 T superconducting solenoid, with data from the gas-ionization muon detectors embedded in the flux-return yoke outside the solenoid. The reconstructed particles are used to build τ lepton candidates, jets, and p_T^{miss} [22–24].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about $4\ \mu\text{s}$ [15]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to a few kHz before data storage [16, 17]. Events are selected using a combination of triggers that require the presence of one, two, or three reconstructed leptons. The trigger efficiency is close to 100% for signal events passing the analysis selection [25].

The simulation of the signal is carried out using the Monte-Carlo (MC) event generator MADGRAPH5_aMC@NLO (MG) version 2.6.5 [5, 6] at NLO accuracy in QCD and in the five-flavor scheme (5FS). The factorization and renormalization scales are estimated dynamically. At leading order (LO) accuracy and in the 5FS, $t\bar{t}WZ$ can be identified through the tree-level partonic process $g b \rightarrow t\bar{t}WZ$. However, at NLO, real corrections of the type $g g \rightarrow t\bar{t}WZb$ arise, that can feature a resonant top quark pair ($t\bar{t}$) in the intermediate state and therefore cause overlap with the LO $t\bar{t}Z$ process [8]. In other words, the emission of an extra b quark implies that the $t\bar{t}WZb$ final state can also occur from the process $g g \rightarrow t\bar{t}Z$, with $t \rightarrow bW^+$ and $\bar{t} \rightarrow \bar{b}W^-$, namely from LO $t\bar{t}Z$ production. The overlap removal is performed using the MADSTR plugin [26], which implements both diagram removal (DR) and diagram subtraction (DS) techniques [27, 28]. For the signal samples, two DR options are used: DR1, which removes from the process amplitude both the squared resonant term and the interference term between resonant and nonresonant contributions, and DR2, which removes only the resonant term. The nominal samples are generated using the DR1 scheme.

The main background processes are simulated with different MC event generators. The dominant backgrounds, $t\bar{t}Z$ and $t\bar{t}Zq$, are generated at NLO accuracy in QCD with MG, while WZ production is simulated using POWHEG v2 [29–33]. Because of larger uncertainties in WZ production in association with a b quark compared to jets from light-flavour quarks and gluons, we split this background into $WZ+b$ and $WZ+j$, respectively. Smaller backgrounds are simulated using MG, POWHEG and MCFM v7.0.1 [34–36]. These comprise di- and tri-boson processes, consolidated under $VV(V)$ where $V = W$ or Z , additional processes with a photon in the final state, $X\gamma$, and processes in which a single top quark or a $t\bar{t}$ pair is produced in association with additional particles, $t(\bar{t})X$, grouping tW , $t\bar{t}H$, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, $t\bar{t}WZ$, and $t\bar{t}ZZ$. The NNPDF 3.1 next-to-NLO (NNLO) parton distribution functions (PDFs) [37] are used for the generation of all samples, together with PYTHIA 8.240 [38] for the modeling of the parton shower (PS) and hadronization. The $t\bar{t}Z$ background cross section is scaled to account for approximate next-to-NNLO (aNNLO) QCD and NLO EW corrections [39]. The underlying event is described using the CP5 tune [40]. Matrix element calculations and PS simulations are matched using the FxFx [6] and MLM [41] algorithms for processes simulated with MG at NLO and LO, respectively. The detector response is simulated with a detailed description of the CMS apparatus based on the GEANT4 toolkit [42]. Additional pp interactions occurring in the same or adjacent bunch crossings (pileup) are modelled using PYTHIA. All simulated samples are produced

taking into account the LHC and detector conditions of the different data-taking periods.

The PF algorithm [21] is used to reconstruct and identify individual particles in the event, combining information from different parts of the detector. The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, identified using the tracking information alone, as described in Section 9.4.1 of Ref. [43]. The energy of electrons is determined from a combination of the electron momentum at the PV, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The momentum of muons is obtained from the curvature of their tracks measured in the tracking system and in the muon detectors. The energy of charged hadrons is evaluated via a combination of their momentum measured in the tracker and the matching of the ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

Electrons and muons must have a transverse momentum $p_T > 10 \text{ GeV}$ for events collected at $\sqrt{s} = 13 \text{ TeV}$, and $p_T > 15 \text{ GeV}$ for events collected at $\sqrt{s} = 13.6 \text{ TeV}$. In addition, electrons must have pseudorapidity $|\eta| < 2.5$ and muons $|\eta| < 2.4$. In this Letter, nonprompt leptons denote reconstructed leptons not originating from the decay of a W or Z boson. To reduce their presence in the final selection, multivariate discriminants are employed [44], obtained from dedicated trainings on 13 TeV [45] and 13.6 TeV [46] data. The lepton identification algorithms for 13 TeV data have efficiencies of 87 and 92% for electrons and muons, respectively. An improved identification is used for 13.6 TeV data, resulting in lower efficiencies of 60 and 84% for electrons and muons, but with a threefold improvement in the rejection of nonprompt leptons.

Jets are reconstructed by clustering the PF candidates using the anti- k_T algorithm [47, 48] with a distance parameter of 0.4. Jets are required to have $p_T > 25 \text{ GeV}$, $|\eta| < 2.5$, and to be separated from selected leptons by a distance $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} > 0.4$. Here, $\Delta\eta$ and $\Delta\phi$ are the differences in pseudorapidity and azimuthal angle between the jet and lepton. Jets are corrected for pileup contributions and detector effects as detailed in Appendix A.1. The DEEPJET algorithm [49–51] is used in 13 TeV data to identify jets that originate from the fragmentation of b quarks (b jets). For 13.6 TeV data, an improved algorithm [52–54] is employed with better performance [55, 56]. We define b jets to be jets that pass a requirement on the algorithm score with an identification efficiency of around 80% for jets truly originating from b quarks and misidentification probabilities of about 1% for light-quark and gluon jets and about 14% for charm quark jets.

We require events in the final selection to have at least three reconstructed leptons, the p_T -leading lepton must have $p_T > 25 \text{ GeV}$ and the p_T -subleading one $p_T > 15 \text{ GeV}$. The events are split into different regions according to the number of leptons, jets, b jets, p_T^{miss} , and invariant mass of the lepton pairs with opposite sign and same flavor (OSSF). The OSSF lepton pairs are considered to lie inside the Z boson mass window if the invariant mass is within $m_Z \pm 15 \text{ GeV}$, where $m_Z = 91.2 \text{ GeV}$ [57]. The signal region (SR) is divided into two parts, depending on the number of leptons: events with three leptons enter the $\text{SR}_{3\ell}$, while events with four leptons are assigned to the $\text{SR}_{4\ell}$. In the $\text{SR}_{3\ell}$, one OSSF lepton pair must be present with mass within the m_Z window. The events must also contain two jets, at least one of which is b tagged. In the $\text{SR}_{4\ell}$, selected events must have exactly one OSSF lepton pair within the m_Z window. Additionally, at least one b-tagged jet must be present in the event.

In addition to the two SRs, we define four control regions (CRs) to constrain the most important backgrounds and validate their modeling. Two CRs are constructed to constrain the ZZ and WZ backgrounds. The other two help to improve the modeling of events containing non-

prompt leptons, originating from Drell–Yan (DY) and $t\bar{t}$ production. Events containing at least one jet and exactly four leptons with two OSSF lepton pairs inside the m_Z window enter the CR_{ZZ} , which is enriched in events from ZZ production. The CR_{WZ} contains events with at least two jets, no b jets, three leptons, two of which are within the m_Z window, and $p_T^{\text{miss}} > 40$ GeV. The CR_{WZ} is also used to improve the description of the jet multiplicity for the WZ process, as detailed in the Appendix A.2. Events entering the CR_{DY} satisfy the same requirements as for the CR_{WZ} , but with an inverted p_T^{miss} selection and no requirement on the number of jets. Finally, the $CR_{t\bar{t}}$ contains events with at least two jets, one of which is b tagged, three leptons but no OSSF lepton pair inside the m_Z window.

The contribution of the nonprompt background is estimated from a fit to data simultaneously with the signal extraction. In the fit, the misidentification rates are left as free parameters and are constrained with the help of distributions in the trailing lepton p_T measured in the CR_{DY} and $CR_{t\bar{t}}$. The trailing lepton is defined as the selected lepton with the smallest p_T . Additional details on the estimation of the nonprompt-lepton background are provided in Appendix A.3.

A transformer ML algorithm based on the particle transformer PART architecture [52] is employed to discriminate between signal and background processes, mainly $t\bar{t}Z$ and WZ production. Four different dedicated models are trained: two for $SR_{3\ell}$ and two for $SR_{4\ell}$, corresponding to trainings with 13 and 13.6 TeV data. In $SR_{3\ell}$, the PART algorithm is set up to have four output nodes, each representing the classification as tWZ, $t\bar{t}Z$, WZ, or any of the minor remaining backgrounds. The events are then divided into three categories based on the maximum value of the output nodes: tWZ, $t\bar{t}Z$, and other backgrounds which combines WZ and minor backgrounds. This results in a region enriched in signal events, a second region mainly used to constrain the $t\bar{t}Z$ background and related uncertainties, and a third region used to constrain other backgrounds. In $SR_{4\ell}$, a binary classifier to distinguish tWZ from background is used due to the lower statistical precision of the data events with four leptons. More details on the algorithm architecture and input variables are provided in Appendix A.4.

The tWZ cross section is measured by maximizing a binned likelihood function in a fit to the data. This function combines the Poisson probabilities of the yields with the predicted signal and background estimates in each bin. It also includes all sources of systematic uncertainties, represented as nuisance parameters, which may influence the observed number of signal or background events. The fit yields the measured signal strength, defined as the ratio of the observed cross section to the expected value, along with its uncertainty. The fit is performed using the CMS statistical analysis program COMBINE [58]. Different distributions enter the fit according to the SR or CR, separately for each center-of-mass energy. Three distributions in the classifier output nodes with five bins each are measured in the $SR_{3\ell}$, corresponding to the values of the tWZ, $t\bar{t}Z$, and other background nodes. Two bins from the classifier output score in the $SR_{4\ell}$ enter the fit. The total number of events measured in the CR_{ZZ} enters the fit to constrain the normalization of the ZZ background. The jet multiplicity distribution is measured in the CR_{WZ} to constrain the uncertainty on the modeling of additional jets produced in association with WZ. The distribution of the trailing lepton p_T is used for the CR_{DY} and $CR_{t\bar{t}}$ to constrain the rates of the nonprompt-lepton identification.

The distributions in the tWZ and $t\bar{t}Z$ output nodes in $SR_{3\ell}$, along with the output of the classifier in $SR_{4\ell}$, are shown in Fig. 2. The distributions are shown for the best fit values of the background normalizations and nuisance parameters corresponding to systematic uncertainties, after a fit to the data.

Several sources of systematic uncertainties are considered, comprising experimental and theoretical components. The experimental ones are the uncertainties in the jet energy corrections




Figure 2: Distributions in the classifier output nodes for the tWZ (upper left) and $t\bar{t}Z$ (upper right) nodes in $SR_{3\ell}$, and in the classifier output node in $SR_{4\ell}$ (lower) after a fit to the data. The dashed band shows the postfit uncertainty. The results from fits to 13 and 13.6 TeV data are combined for illustration purposes. The ratio of data to the predictions from MC simulation is shown below the distributions.

and resolution [23], unclustered energy [24], the description of pileup in the simulation [59], the trigger [15, 16], b tagging [49], and lepton efficiency corrections [18, 19]. The integrated luminosity has an uncertainty of 1.6% for the 2016–2018 data-taking years [60–62], and 1.4 and 1.3% for 2022 and 2023, respectively [63, 64]. The theoretical uncertainties include uncertainties in the PDFs, the PS simulation, and the matrix element renormalization and factorization scales. Additional sources of uncertainty affecting the signal are taken into account by considering differences between the DR1 and DR2 subtraction schemes. Variations from DS are not considered, as these are smaller than the ones derived with DR2. The uncertainty associated with the modeling of additional jets in WZ production is also included, more details are given in the Appendix A.2. In addition, theoretical uncertainties in the cross sections of the background processes are considered. A normalization uncertainty of 7% is employed for the $t\bar{t}Z$ cross section [65, 66]. Two different normalization uncertainties are considered for the WZ and ZZ processes, where we assign a 20% uncertainty on VV+b production [25] and a 10% uncertainty on VV+j [67]. An uncertainty of 11% is used for tZq production [25], and a 10% uncertainty for VV(V) [68, 69] and $X\gamma$ [70, 71]. Uncertainties of 5 and 10% are applied to the $t\bar{t}$ [72] and DY processes [73]. A 25% uncertainty is considered for the $t\bar{t}W$ process where the measurements do not agree with the SM predictions by about this amount [74], and a 20% uncertainty is assigned to $t(\bar{t})X$ [45, 75, 76]. Additional uncertainties describe the rate of misidentified non-prompt leptons as detailed in Appendix A.3. Finally, uncertainties arising from the finite size of simulated samples are accounted for using the Barlow–Beeston light method [77].

The measured signal strength of tWZ production is $\mu_{tWZ} = 1.77 \pm 0.23$ (stat) ± 0.22 (syst) with an observed statistical significance of 5.8 s.d. compared to the background-only hypothesis, with 3.5 s.d. expected. The theoretical predictions come with an uncertainty on the signal strength of 0.06, thus the result is 2.3 s.d. larger than the SM expectation, consistent with the previous measurement [11]. The signal strength using 13 TeV data only is 1.82 ± 0.28 (stat) ± 0.26 (syst), in agreement with the one from 13.6 TeV data only, 1.64 ± 0.42 (stat) ± 0.31 (syst). The corresponding observed significances are 5.1 s.d. and 3.4 s.d. for 13 and 13.6 TeV data, with expected significances of 3.0 s.d. and 2.1 s.d., respectively. The measured cross sections for tWZ production are 248 ± 38 (stat) ± 35 (syst) fb and 242 ± 62 (stat) ± 46 (syst) fb for $\sqrt{s} = 13$ and 13.6 TeV, respectively. Signal strengths measured in the different data-taking periods and leptonic channels are presented in Appendix A.5. The dominant systematic uncertainty is the normalization of the $t\bar{t}Z$ process with a contribution of 14% to the total uncertainty. Other relevant uncertainties are the normalization of the WZ+b process and the uncertainties in the jet energy corrections, with contributions of 9 and 6%, respectively. We determine the signal strengths of tWZ and $t\bar{t}Z$ production in a simultaneous fit, without external constraints on the $t\bar{t}Z$ cross section. The result of this fit is shown in Fig. 3. The measured signal strengths are $\mu_{tWZ} = 1.88 \pm 0.30$ (stat) ± 0.15 (syst) and $\mu_{t\bar{t}Z} = 0.91 \pm 0.04$ (stat) ± 0.03 (syst), also resulting in a significance of more than five s.d. for tWZ production. The measured $t\bar{t}Z$ signal strength is compatible within two s.d. with the most recent theoretical prediction at aN^3LO QCD and NLO EW [39]. It is compatible with previous results on $t\bar{t}Z$ production [65, 66, 78] when considering the 12% increase in the predicted cross section compared to that used in these measurements. The correlation between μ_{tWZ} and $\mu_{t\bar{t}Z}$ is reduced compared to the previous CMS analysis [11] because of the improved ML algorithm and the separation of the $SR_{3\ell}$ into three categories. The combined effect is a better constraint on μ_{tWZ} largely independent of the value of $\mu_{t\bar{t}Z}$.

In summary, a measurement of tWZ production in proton-proton (pp) collisions has been presented in events with three and four charged leptons. The analysis is performed using data recorded with the CMS detector at center-of-mass energies of 13 and 13.6 TeV, corresponding to a total integrated luminosity of 200 fb^{-1} . Advanced machine-learning algorithms are




Figure 3: Likelihood scan of the signal strengths for tWZ and $t\bar{t}Z$ production. The black cross shows the best fit value, while the black diamond indicates the SM expected value. The contours in different shades of blue correspond to the 68, 95, and 99% confidence level.

employed to significantly improve the sensitivity of the analysis compared to an earlier analysis. Analyzing the same data, an expected significance of 3.0 standard deviations (s.d.) is achieved, compared to 1.4 s.d. obtained previously. Combining the 13 and 13.6 TeV data, a significance of 5.8 s.d. is observed. The measured signal strength $\mu_{tWZ} = 1.77 \pm 0.32$ (tot) agrees with the standard model prediction within 2.3 s.d. The measured cross sections are 248 ± 52 fb and 242 ± 77 fb for $\sqrt{s} = 13$ and 13.6 TeV, respectively. This is the first observation of tWZ production in pp collisions. This measurement represents a significant step forward in our understanding of rare processes involving the top quark and electroweak bosons. It opens up the possibility for future studies of the top quark electroweak interactions in the context of the standard model and its extensions.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan);

MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

References

- [1] S. Rappoccio, “The experimental status of direct searches for exotic physics beyond the standard model at the Large Hadron Collider”, *Rev. Phys.* **4** (2019) 100027, doi:10.1016/j.revip.2018.100027, arXiv:1810.10579.
- [2] ATLAS Collaboration, “Exploration at the high-energy frontier: ATLAS Run 2 searches investigating the exotic jungle beyond the standard model”, *Phys. Rept.* **1116** (2025) 301, doi:10.1016/j.physrep.2024.10.001, arXiv:2403.09292.
- [3] CMS Collaboration, “Stairway to discovery: A report on the CMS programme of cross section measurements from millibarns to femtobarns”, *Phys. Rept.* **1115** (2025) 3, doi:10.1016/j.physrep.2024.11.005, arXiv:2405.18661.
- [4] D. London and J. Matias, “*B* flavour anomalies: 2021 theoretical status report”, *Ann. Rev. Nucl. Part. Sci.* **72** (2022) 37, doi:10.1146/annurev-nucl-102020-090209, arXiv:2110.13270.
- [5] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [6] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [7] F. Maltoni, L. Mantani, and K. Mimasu, “Top-quark electroweak interactions at high energy”, *JHEP* **10** (2019) 004, doi:10.1007/JHEP10(2019)004, arXiv:1904.05637.
- [8] H. El Faham, F. Maltoni, K. Mimasu, and M. Zaro, “Single top production in association with a WZ pair at the LHC in the SMEFT”, *JHEP* **01** (2022) 100, doi:10.1007/JHEP01(2022)100, arXiv:2111.03080.
- [9] J. Keaveney, “Constraining the SMEFT with a differential cross section measurement of tWZ production at the HL-LHC”, *Phys. Rev. D* **107** (2023) 036021, doi:10.1103/PhysRevD.107.036021, arXiv:2107.01053.
- [10] A. Belvedere, C. Englert, R. Kogler, and M. Spannowsky, “Dispelling the $\sqrt{\mathcal{L}}$ myth for the High-Luminosity LHC”, *Eur. Phys. J. C* **84** (2024) 715, doi:10.1140/epjc/s10052-024-13032-w, arXiv:2402.07985.
- [11] CMS Collaboration, “Evidence for tWZ production in proton-proton collisions at $\sqrt{s} = 13$ TeV in multilepton final states”, *Phys. Lett. B* **855** (2024) 138815, doi:10.1016/j.physletb.2024.138815, arXiv:2312.11668.
- [12] “HEPData record for this analysis”, 2025. doi:10.17182/hepdata.158454.v1.
- [13] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.

- [14] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [15] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [16] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [17] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [18] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [19] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [20] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [21] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [22] CMS Collaboration, “Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P10005, doi:10.1088/1748-0221/13/10/P10005, arXiv:1809.02816.
- [23] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [24] CMS Collaboration, “Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector”, *JINST* **14** (2019) P07004, doi:10.1088/1748-0221/14/07/P07004, arXiv:1903.06078.
- [25] CMS Collaboration, “Inclusive and differential cross section measurements of single top quark production in association with a Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **02** (2022) 107, doi:10.1007/JHEP02(2022)107, arXiv:2111.02860.
- [26] S. Frixione et al., “Automated simulations beyond the standard model: Supersymmetry”, *JHEP* **12** (2019) 008, doi:10.1007/JHEP12(2019)008, arXiv:1907.04898.
- [27] S. Frixione et al., “Single-top hadroproduction in association with a W boson”, *JHEP* **07** (2008) 029, doi:10.1088/1126-6708/2008/07/029, arXiv:0805.3067.

-
- [28] C. D. White, S. Frixione, E. Laenen, and F. Maltoni, “Isolating Wt production at the LHC”, *JHEP* **11** (2009) 074, doi:10.1088/1126-6708/2009/11/074, arXiv:0908.0631.
- [29] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [30] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: The POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [31] S. Frixione, G. Ridolfi, and P. Nason, “A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction”, *JHEP* **09** (2007) 126, doi:10.1088/1126-6708/2007/09/126, arXiv:0707.3088.
- [32] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: The POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [33] E. Re, “Single-top Wt -channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [34] J. M. Campbell and R. K. Ellis, “An update on vector boson pair production at hadron colliders”, *Phys. Rev. D* **60** (1999) 113006, doi:10.1103/PhysRevD.60.113006, arXiv:hep-ph/9905386.
- [35] J. M. Campbell, R. K. Ellis, and C. Williams, “Vector boson pair production at the LHC”, *JHEP* **07** (2011) 018, doi:10.1007/JHEP07(2011)018, arXiv:1105.0020.
- [36] J. M. Campbell, R. K. Ellis, and W. T. Giele, “A multi-threaded version of MCFM”, *Eur. Phys. J. C* **75** (2015) 246, doi:10.1140/epjc/s10052-015-3461-2, arXiv:1503.06182.
- [37] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
- [38] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [39] N. Kidonakis and C. Foster, “Higher-order soft-gluon corrections for $t\bar{t}Z$ cross sections”, *Phys. Lett. B* **860** (2025) 139146, doi:10.1016/j.physletb.2024.139146, arXiv:2410.01214.
- [40] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA 8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [41] J. Alwall et al., “Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions”, *Eur. Phys. J. C* **53** (2008) 473, doi:10.1140/epjc/s10052-007-0490-5, arXiv:0706.2569.
- [42] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.

- [43] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
doi:10.17181/CERN.VU8I.D59J.
- [44] CMS Collaboration, “Muon identification using multivariate techniques in the cms experiment in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **19** (2024) P02031,
doi:10.1088/1748-0221/19/02/P02031, arXiv:2310.03844.
- [45] CMS Collaboration, “Observation of four top quark production in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **847** (2023) 138290,
doi:10.1016/j.physletb.2023.138290, arXiv:2305.13439.
- [46] CMS Collaboration, “Measurement of the inclusive WZ production cross section in pp collisions at $\sqrt{s} = 13.6$ TeV”, *JHEP* **04** (2025) 115, doi:10.1007/JHEP04(2025)115,
arXiv:2412.02477.
- [47] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [48] M. Cacciari, G. P. Salam, and G. Soyez, “FASTJET user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [49] CMS Collaboration, “Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV”, *JINST* **13** (2018) P05011,
doi:10.1088/1748-0221/13/05/P05011, arXiv:1712.07158.
- [50] E. Bols et al., “Jet flavour classification using DeepJet”, *JINST* **15** (2020) P12012,
doi:10.1088/1748-0221/15/12/P12012, arXiv:2008.10519.
- [51] CMS Collaboration, “Performance summary of AK4 jet b tagging with data from proton-proton collisions at 13 TeV with the CMS detector”, CMS Detector Performance Note CMS-DP-2023-005, 2023.
- [52] H. Qu, C. Li, and S. Qian, “Particle transformer for jet tagging”, in *Proc. 39th International Conference on Machine Learning (ICML 2022): Baltimore MD, USA, July 17–23. 2022.*
arXiv:2202.03772.
- [53] CMS Collaboration, “Transformer models for heavy flavor jet identification”, CMS Detector Performance Note CMS-DP-2022-050, 2022.
- [54] CMS Collaboration, “Adversarial training for b-tagging algorithms in CMS”, CMS Detector Performance Note CMS-DP-2022-049, 2022.
- [55] CMS Collaboration, “Run 3 commissioning results of heavy-flavor jet tagging at $\sqrt{s} = 13.6$ TeV with CMS data using a modern framework for data processing”, CMS Detector Performance Note CMS-DP-2022-024, 2024.
- [56] CMS Collaboration, “Performance summary of AK4 jet b tagging with data from 2022 proton-proton collisions at 13.6 TeV with the CMS detector”, CMS Detector Performance Note CMS-DP-2022-025, 2024.
- [57] Particle Data Group, “Review of particle physics”, *Phys. Rev. D* **110** (2024) 030001,
doi:10.1103/PhysRevD.110.030001.

-
- [58] CMS Collaboration, “The CMS statistical analysis and combination tool: COMBINE”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [59] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [60] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [61] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [62] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [63] CMS Collaboration, “Luminosity measurement in proton-proton collisions at 13.6 TeV in 2022 at CMS”, CMS Physics Analysis Summary CMS-PAS-LUM-22-001, 2024.
- [64] CMS Collaboration, “Measurement of the offline integrated luminosity for the CMS proton-proton collision dataset recorded in 2023”, CMS Detector Performance Note CMS-DP-2024-068, 2024.
- [65] CMS Collaboration, “Measurements of inclusive and differential cross sections for top quark production in association with a Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **02** (2025) 177, doi:10.1007/JHEP02(2025)177, arXiv:2410.23475.
- [66] ATLAS Collaboration, “Inclusive and differential cross-section measurements of $t\bar{t}Z$ production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, including EFT and spin-correlation interpretations”, *JHEP* **07** (2024) 163, doi:10.1007/JHEP07(2024)163, arXiv:2312.04450.
- [67] CMS Collaboration, “Measurement of the inclusive and differential WZ production cross sections, polarization angles, and triple gauge couplings in pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2022) 032, doi:10.1007/JHEP07(2022)032, arXiv:2110.11231.
- [68] CMS Collaboration, “Observation of the production of three massive gauge bosons at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **125** (2020) 151802, doi:10.1103/PhysRevLett.125.151802, arXiv:2006.11191.
- [69] ATLAS Collaboration, “Observation of $W\gamma\gamma$ triboson production in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett. B* **848** (2024) 138400, doi:10.1016/j.physletb.2023.138400, arXiv:2308.03041.
- [70] ATLAS Collaboration, “Observation of $WZ\gamma$ production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **132** (2024) 021802, doi:10.1103/PhysRevLett.132.021802, arXiv:2305.16994.
- [71] CMS Collaboration, “Observation of $WW\gamma$ production and search for $H\gamma$ production in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **132** (2024) 121901, doi:10.1103/PhysRevLett.132.121901, arXiv:2310.05164.

- [72] M. Czakon and A. Mitov, “Top++: A program for the calculation of the top-pair cross-section at hadron colliders”, *Comput. Phys. Commun.* **185** (2014) 2930, doi:10.1016/j.cpc.2014.06.021, arXiv:1112.5675.
- [73] CMS Collaboration, “Measurement of differential cross sections for the production of a Z boson in association with jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **108** (2023) 052004, doi:10.1103/PhysRevD.108.052004, arXiv:2205.02872.
- [74] CMS Collaboration, “Measurement of the cross section of top quark-antiquark pair production in association with a W boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2023) 219, doi:10.1007/JHEP07(2023)219, arXiv:2208.06485.
- [75] CMS Collaboration, “Measurement of inclusive and differential cross sections for single top quark production in association with a W boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2023) 046, doi:10.1007/JHEP07(2023)046, arXiv:2208.00924.
- [76] CMS Collaboration, “A portrait of the Higgs boson by the CMS experiment ten years after the discovery”, *Nature* **607** (2022) 60, doi:10.1038/s41586-022-04892-x, arXiv:2207.00043. [Erratum: doi:10.1038/s41586-023-06164-8].
- [77] R. Barlow and C. Beeston, “Fitting using finite Monte Carlo samples”, *Comput. Phys. Commun.* **77** (1993) 219, doi:10.1016/0010-4655(93)90005-W.
- [78] CMS Collaboration, “Measurement of top quark pair production in association with a Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **03** (2020) 056, doi:10.1007/JHEP03(2020)056, arXiv:1907.11270.
- [79] D. Bertolini, P. Harris, M. Low, and N. Tran, “Pileup per particle identification”, *JHEP* **10** (2014) 59, doi:10.1007/JHEP10(2014)059, arXiv:1407.6013.
- [80] CMS Collaboration, “Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s} = 8$ TeV”, *JINST* **10** (2015) P02006, doi:10.1088/1748-0221/10/02/P02006, arXiv:1411.0511.

A Additional material

A.1 Pileup mitigation

Pileup can contribute additional tracks and calorimetric energy depositions to the jet momentum. For data from 2016–2018, we mitigate this effect by discarding charged particles identified to be originating from pileup vertices and apply an offset correction to correct for remaining pileup contributions [21, 59]. For 2022–2023 data, we apply the pileup-per-particle identification (PUPPI) algorithm [59, 79], weighting all PF candidates by their probability to originate from the PV [24]. Jet energy corrections are derived from data and simulation to bring the measured response of jets to that of particle level jets on average [23]. The missing transverse momentum vector $\vec{p}_T^{\text{miss}}$ is the negative vector sum of the transverse momenta of all PF candidates [24, 80]. Corrections to the jet energies are propagated to $\vec{p}_T^{\text{miss}}$.

A.2 Jet multiplicity reweighting

The fully leptonic decay of the WZ process involves at most one jet from the matrix element calculation at NLO in QCD. Thus, additional jet emissions are primarily simulated by the PS, which may lead to mis-modeling in the shape of the jet multiplicity distribution. This can be particularly relevant when selections that are not inclusive in the number of jets are applied. To improve the modeling of the jet multiplicity, a simulation-to-data reweighting is performed in the CR_{WZ} . Simulated contributions from processes other than WZ are subtracted from data, and a correction is derived as the ratio between the observed data yield and the WZ simulation as a function of the jet multiplicity. A shape uncertainty is assigned to this correction by considering two alternative configurations: one where the correction factor is doubled and another where no correction is applied.




Figure A.1: Distribution in the p_T of the 3rd leading lepton for the CR_{DY} (left) and CR_{tt} (right) after a fit to data. The dashed band shows the postfit uncertainty. The results from fits to 13 and 13.6 TeV data are combined for illustration purposes. The ratio of data to the predictions from MC simulations is shown below the distributions.

A.3 Background from nonprompt leptons

The nonprompt-lepton contribution is estimated from a fit to data using simulated templates based on the origin of the misidentified leptons. Events containing at least one nonprompt lepton are divided into three categories: nonprompt leptons from photons, b jets, and light-quark and gluon jets. For 13.6 TeV data, nonprompt leptons from b-quark, light-quark, and gluon jets are combined into a single category because of the smaller number of events from nonprompt leptons due to the improved lepton-identification algorithm. We introduce unconstrained parameters that describe the lepton misidentification rates as functions of p_T . For nonprompt leptons from photons, these parameters are considered to be normal-distributed around the MC predictions, with prior uncertainties of 50%. Uncertainties accounting for shape differences in the trailing-lepton p_T distributions between nonprompt leptons of different flavors within the same category are considered as an additional uncertainty. We find that these have a negligible impact on the analysis. In the fit, the two distributions in the trailing-lepton p_T measured in the CR_{DY} and $\text{CR}_{\text{t}\bar{\text{t}}}$ are used. These are shown in Fig. A.1 after the fit to data. The nonprompt backgrounds constitute about one-third of all selected events in these two regions. The trailing-lepton p_T distributions show sensitivity to the origin of the nonprompt lepton and can thus constrain the parameters describing the misidentification rates. These are measured independently for each data-taking period. The largest correction factors are measured for nonprompt leptons from light-quark and gluon jets, resulting in a nonprompt-lepton background that is 40–70% lower than the simulation-based prediction, depending on the data-taking period. The associated uncertainties are around 50%, due to the small statistical power of this category. The correction factors associated with the nonprompt leptons from b jets are closer to unity, where the largest correction shifts the prediction up by 40%, with uncertainties of about 20%.

A.4 The machine-learning algorithm

The PART algorithm takes as inputs two different set of variables: *Particles* and *Interactions*. The *Particles* input corresponds to a list of individual physics objects where each particle is described by a set of features. It is divided into two categories: *leptons* and *jets*. The components of the four-momentum of each lepton and jet in every event are used as inputs, as well as the $\vec{p}_T^{\text{miss}}$. Additionally, the following set of features is considered: $\log(p_T/\text{GeV})$, $\log(E/\text{GeV})$, η , ϕ , b-tagging information (only for jets), and a number that encodes the flavor of the lepton. The agreement between data and simulation has been verified for distributions in all input variables. The *interactions* encode correlations between pairs of particles considering the following variables: $\log(\Delta)$, $\log(k_T/\text{GeV})$, $\log(z)$, and $\log(m^2/\text{GeV}^2)$. These are defined between two particles a and b as

$$\begin{aligned}\Delta &= \sqrt{(y_a - y_b)^2 + (\phi_a - \phi_b)^2}, \\ k_T &= \min(p_{T,a}, p_{T,b})\Delta, \\ z &= \min(p_{T,a}, p_{T,b}) / (p_{T,a} + p_{T,b}), \\ m^2 &= (E_a + E_b)^2 - |\mathbf{p}_a + \mathbf{p}_b|^2,\end{aligned}$$

where y , ϕ , and $\mathbf{p}$ are the rapidity, azimuthal angle, and the three-momentum. The *interactions* allow for each particle to incorporate information from all the other particles using both the particle features and their pairwise relationships, enabling the network to capture both local and global event characteristics. To avoid overtraining, two versions of the PART algorithm are trained for each signal region and center-of-mass energy. An algorithm trained on one half of the simulated data is employed on the other half during the analysis. In this way, each algorithm is trained on and applied to separate halves of the simulated samples. The agreement




Figure A.2: Values of the signal strengths for tWZ production obtained from the different data-taking periods (left), and from events with different lepton flavors (right): three electrons (eee), two electrons and one muon ($ee\mu$), one electron and two muons ($e\mu\mu$), and three muons ($\mu\mu\mu$). If there are more than three leptons in an event, the three p_T -leading leptons are used to determine the flavor. The combined results from 13 TeV data (Run 2) and from 13.6 TeV data (Run 3) are also shown. The result from the combined fit to 13 and 13.6 TeV data is shown as a red line with the corresponding uncertainty shown as a red band. The SM prediction is shown as a gray band.

between data and simulation has also been verified in the $\text{CR}_{t\bar{t}}$ for the algorithms used in the $\text{SR}_{3\ell}$, and in the CR_{ZZ} for the algorithms employed in the $\text{SR}_{4\ell}$. These CRs have the same number of physics objects as the corresponding SRs, and the data distributions in the output of the algorithms show good agreement to the simulated ones.

A.5 Individual results

The signal strength is also measured for the different data-taking periods and leptonic channels, defined by the flavor of the three leptons with largest p_T in the event. Comparisons of the obtained signal strengths are shown in Figure A.2. We find good agreement between the measured signal strengths in the different data-taking periods and among the signal strengths from different lepton flavors.

B The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , M. Dragicevic , C. Giordano, P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , M. Matthewman, I. Mikulec , J. Schieck² , R. Schöfbeck² , D. Schwarz , M. Shooshtari , M. Sonawane , W. Waltenberger , C.-E. Wulz² 

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , D. Ocampo Henao , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

J. Bierkens , N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , P. Kashko , S. Lowette , I. Makarenko , D. Müller , J. Song , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , D. Marckx , K. Skovpen , N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , A. Giammanco , A.O. Guzel , V. Lemaitre, J. Lidrych , P. Malek , P. Mastrapasqua , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Barroso Ferreira Filho , E. Coelho , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote⁵ , A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶ , M. Costa Reis , E.M. Da Costa , G.G. Da Silva⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza , S. S. Jesus , T. Laux Kuhn⁷ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁸ 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , F. Damas , T.R. Fernandez Perez Tomei , E.M. Gregores , B. Lopes Da Costa , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , P. Danev, R. Hadjiiska , P. Iaydjiev , M. Shopova , G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Universidad Tecnica Federico Santa Maria, Valparaiso, Chile

W. Brooks 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Wang , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, X. Wang 

Institute of High Energy Physics, Beijing, China

G.M. Chen⁹ , H.S. Chen⁹ , M. Chen⁹ , Y. Chen , Q. Hou , X. Hou, F. Iemmi , C.H. Jiang, A. Kapoor¹⁰ , H. Liao , G. Liu , Z.-A. Liu¹¹ , J.N. Song¹¹, S. Song, J. Tao , C. Wang⁹, J. Wang , H. Zhang , J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, Q. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, C. Quaranta , X. Sun , D. Wang , J. Wang, H. Yang, M. Zhang, Y. Zhao, C. Zhou 

State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer^{12,13}, Z. Cui¹³, B. Li¹⁴, H. Wang , K. Yi¹⁵ , J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao¹⁶ 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , C. Nicolaou, L. Paizanos , F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger[†] , M. Finger Jr. 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

H. Abdalla¹⁷ , Y. Assran^{18,19}

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad , A. Hussein, H. Mohammed 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , N. Seeba , L. Tani 

Department of Physics, University of Helsinki, Helsinki, Finland

A. Milieva , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , R. Kumar Verma , T. Lampén , K. Lassila-Perini , B. Lehtela , S. Lehti , T. Lindén , N.R. Mancilla Xinto , M. Myllymäki , M.m. Rantanen , S. Saariokari , N.T. Toikka , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

H. Kirschenmann , P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, P. Devouge, J.L. Faure , F. Ferri , P. Gagne, S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , S. Ronchi , M.Ö. Sahin , A. Savoy-Navarro²⁰ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , O. Davignon , A. De Wit , T. Debnath , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , L. Kalipoliti , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²¹ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard 

G. Coulon, S. Falke^{ID}, U. Goerlach^{ID}, R. Haeberle^{ID}, A.-C. Le Bihan^{ID}, M. Meena^{ID}, O. Poncet^{ID}, G. Saha^{ID}, P. Vaucelle^{ID}

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio^{ID}

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron^{ID}, B. Blancon^{ID}, G. Boudoul^{ID}, N. Chanon^{ID}, D. Contardo^{ID}, P. Depasse^{ID}, H. El Mamouni, J. Fay^{ID}, S. Gascon^{ID}, M. Gouzevitch^{ID}, C. Greenberg^{ID}, G. Grenier^{ID}, B. Ille^{ID}, E. Jourdhuy, I.B. Laktineh, M. Lethuillier^{ID}, B. Massoteau^{ID}, L. Mirabito, A. Purohit^{ID}, M. Vander Donckt^{ID}, J. Xiao^{ID}

Georgian Technical University, Tbilisi, Georgia

A. Khvedelidze²²^{ID}, I. Lomidze^{ID}, Z. Tsamalaidze²²^{ID}

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta^{ID}, S. Consuegra Rodríguez^{ID}, L. Feld^{ID}, K. Klein^{ID}, M. Lipinski^{ID}, D. Meuser^{ID}, P. Natland^{ID}, V. Oppenländer, A. Pauls^{ID}, D. Pérez Adán^{ID}, N. Röwert^{ID}, M. Teroerde^{ID}

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Daumann, S. Diekmann^{ID}, A. Dodonova^{ID}, N. Eich^{ID}, D. Eliseev^{ID}, F. Engelke^{ID}, J. Erdmann^{ID}, M. Erdmann^{ID}, B. Fischer^{ID}, T. Hebbeker^{ID}, K. Hoepfner^{ID}, F. Ivone^{ID}, A. Jung^{ID}, N. Kumar^{ID}, M.y. Lee^{ID}, F. Mausolf^{ID}, M. Merschmeyer^{ID}, A. Meyer^{ID}, F. Nowotny, A. Pozdnyakov^{ID}, W. Redjeb^{ID}, H. Reithler^{ID}, U. Sarkar^{ID}, V. Sarkisovi^{ID}, A. Schmidt^{ID}, C. Seth, A. Sharma^{ID}, J.L. Spah^{ID}, V. Vaulin, S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M.R. Beckers^{ID}, C. Dziwok^{ID}, G. Flügge^{ID}, N. Hoeflich^{ID}, T. Kress^{ID}, A. Nowack^{ID}, O. Pooth^{ID}, A. Stahl^{ID}, A. Zotz^{ID}

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen^{ID}, A. Abel, M. Aldaya Martin^{ID}, J. Alimena^{ID}, S. Amoroso, Y. An^{ID}, I. Andreev^{ID}, J. Bach^{ID}, S. Baxter^{ID}, M. Bayatmakou^{ID}, H. Becerril Gonzalez^{ID}, O. Behnke^{ID}, A. Belvedere^{ID}, F. Blekman²³^{ID}, K. Borras²⁴^{ID}, A. Campbell^{ID}, S. Chatterjee^{ID}, L.X. Coll Saravia^{ID}, G. Eckerlin, D. Eckstein^{ID}, E. Gallo²³^{ID}, A. Geiser^{ID}, V. Guglielmi^{ID}, M. Guthoff^{ID}, A. Hinzmann^{ID}, L. Jeppe^{ID}, M. Kasemann^{ID}, C. Kleinwort^{ID}, R. Kogler^{ID}, M. Komm^{ID}, D. Krücker^{ID}, W. Lange, D. Leyva Pernia^{ID}, K.-Y. Lin^{ID}, K. Lipka²⁵^{ID}, W. Lohmann²⁶^{ID}, J. Malvaso^{ID}, R. Mankel^{ID}, I.-A. Melzer-Pellmann^{ID}, M. Mendizabal Morentin^{ID}, A.B. Meyer^{ID}, G. Milella^{ID}, K. Moral Figueroa^{ID}, A. Mussgiller^{ID}, L.P. Nair^{ID}, J. Niedziela^{ID}, A. Nürnberg^{ID}, J. Park^{ID}, E. Ranken^{ID}, A. Raspereza^{ID}, D. Rastorguev^{ID}, L. Rygaard^{ID}, M. Scham^{27,24}^{ID}, S. Schnake²⁴^{ID}, P. Schütze^{ID}, C. Schwanenberger²³^{ID}, D. Selivanova^{ID}, K. Sharko^{ID}, M. Shchedrolosiev^{ID}, D. Stafford^{ID}, M. Torkian, F. Vazzoler^{ID}, A. Ventura Barroso^{ID}, R. Walsh^{ID}, D. Wang^{ID}, Q. Wang^{ID}, K. Wichmann, L. Wiens²⁴^{ID}, C. Wissing^{ID}, Y. Yang^{ID}, S. Zakharov, A. Zimmermann Castro Santos^{ID}

University of Hamburg, Hamburg, Germany

A.R. Alves Andrade^{ID}, M. Antonello^{ID}, S. Bollweg, M. Bonanomi^{ID}, K. El Morabit^{ID}, Y. Fischer^{ID}, M. Frahm, E. Garutti^{ID}, A. Grohsjean^{ID}, A.A. Guvenli^{ID}, J. Haller^{ID}, D. Hundhausen, G. Kasieczka^{ID}, P. Keicher^{ID}, R. Klanner^{ID}, W. Korcari^{ID}, T. Kramer^{ID}, C.c. Kuo, F. Labe^{ID}, J. Lange^{ID}, A. Lobanov^{ID}, L. Moureaux^{ID}, A. Nigamova^{ID}, K. Nikolopoulos^{ID}, A. Paasch^{ID}, K.J. Pena Rodriguez^{ID}, N. Prouvost, T. Quadfasel^{ID}

B. Raciti , M. Rieger , D. Savoie , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , R. Ward , B. Wiederspan, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁸ , M. Horzela , F. Hummer , U. Husemann , J. Kieseler , M. Klute , R. Kunnilan Muhammed Rafeek, O. Lavoryk , J.M. Lawhorn , A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa , L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , C. Verstege , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , R. Wolf , W.D. Zeuner , X. Zuo

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou , G. Daskalakis , A. Kyriakis 

National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , E. Tzovara , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

T. Chatzistavrou , G. Karapostoli , K. Kousouris , E. Siamarkou, G. Tsipolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

D. Druzhkin , C. Hajdu , D. Horvath^{29,30} , K. Márton, A.J. Rádl³¹ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³² , M.M.A. Gadallah³³ , Á. Kadlecik , M. León Coello , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³² , F. Nemes³² , T. Novak , I. Szanyi³⁴ 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁵ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , S. Phor , K. Ranjan , M.K. Saini

University of Hyderabad, Hyderabad, India

S. Acharya³⁶ , B. Gomber , B. Sahu³⁶ 

Indian Institute of Technology Kanpur, Kanpur, India

S. Mukherjee 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁷ , T. Mishra , P.R. Pujahari , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

IISER Mohali, India, Mohali, India

B.K. Sirasva

Tata Institute of Fundamental Research-A, Mumbai, India

L. Bhatt, S. Dugad , G.B. Mohanty , M. Shelake , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , S. Barman³⁸ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , M. Maity³⁸, G. Majumder , K. Mazumdar , S. Parolia , R. Saxena , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁹ , D. Maity⁴⁰ , P. Mal , K. Naskar⁴⁰ , A. Nayak⁴⁰ , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese⁴⁰ , D. Vats⁴⁰ 

Indian Institute of Science Education and Research (IISER), Pune, India

A. Alpana , S. Dube , P. Hazarika , B. Kansal , A. Laha , R. Sharma , S. Sharma , K.Y. Vaish 

Indian Institute of Technology Hyderabad, Telangana, India

S. Ghosh 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴¹ , A. Jafari⁴² , V. Sedighzadeh Dalavi , M. Zeinali⁴³ 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri , S. Chenarani⁴⁴ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁵ 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , M. Barbieri^{a,b}, M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,46} , N. Ferrara^{a,c} , L. Fiore^a , L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , F. Nenna^{a,b} , S. Nuzzo^{a,b} , A. Pellicchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} 

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , P. Capiluppi^{a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,47} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarra^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,48} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,48} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

J. Altork^{a,b} , P. Assiouras^a , G. Barbagli^a , G. Bardelli^a , M. Bartolini^{a,b} , A. Calandri^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , L. Damenti^{a,b} , E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁹ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , V. Camagni^{a,b} , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.R. Kim^a , G. Lavizzari^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , C. Di Fraia^{a,b} , F. Fabozzi^{a,c} , L. Favilla^{a,d} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,50} , P. Paolucci^{a,28} , B. Rossi^a 

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

P. Azzi^a , N. Bacchetta^{a,51} , M. Benettoni^a , D. Bisello^{a,b} , P. Bortignon^{a,c} , G. Bortolato^{a,b} , A.C.M. Bulla^{a,c} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^{a,52} , U. Gasparini^{a,b} , S. Giorgetti^a , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , P. Montagna^{a,b} , M. Pelliccioni^a , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Della Penna^{a,b} , L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,53} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy;

Università di Siena^d, Siena, Italy

C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , L. Calligaris^{a,b} , R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,d} , M. Cipriani^{a,b} , R. Dell'Orso^a , S. Donato^{a,b} , R. Forti^{a,b} , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^{a,b} , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b} , S. Nandan^a , F. Palla^a , M. Riggirello^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,54} , T. Sarkar^a , A. Scribano^a , P. Solanki^{a,b} , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , S.C. Behera^a , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , F. De Ruggi^{a,b} , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , F. Errico^a , L. Frosina^{a,b} , R. Gargiulo^{a,b} , B. Harikrishnan^{a,b} , F. Lombardi^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , R. Paramatti^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , L. Lanteri^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Ruspà^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , E. Vlasov^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} , R. Delli Gatti^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, T. Kim , D. Lee, H. Lee , J. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon , J. Seo 

Hanyang University, Seoul, Korea

E. Asilar , F. Carnevali , J. Choi⁵⁵ , T.J. Kim , Y. Ryou 

Korea University, Seoul, Korea

S. Ha , S. Han, B. Hong , J. Kim , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , J. Shin , S. Yang 

Sejong University, Seoul, Korea

Y. Kang , H. S. Kim , Y. Kim , S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , H. Kim , J. Kim , T. Kim, Y. Kim, Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , J. Shin , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, D. Kim , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

G. Cho, K. Hwang , B. Kim , S. Kim, K. Lee , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , O.M. Eberlins , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs , R. Plese , A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , S. Nargelas , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁶ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , A. Cota Rodriguez , L.E. Cuevas Picos, H.A. Encinas Acosta, L.G. Gallegos Maríñez, J.A. Murillo Quijada , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma , R. Lopez-Fernandez , J. Mejia Guisao , R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar , A. Awais , M.I.M. Awan, W.A. Khan 

AGH University of Krakow, Krakow, Poland

V. Avati, L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski 

National Centre for Nuclear Research, Swierk, Poland

M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , L. Markovic , P. Milenovic , V. Milosevic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic, M. Stojanovic 

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Alcalde Martinez , J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , J. Cuchillo Ortega, B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , M. Gonzalez Hernandez , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , R. Paz Herrera , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Ayllon Torresano , A. Cardini , J. Cuevas , J. Del Riego Badas , D. Estrada Acevedo , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon , J. Prado Pico , A. Soto Rodríguez , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasiosa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Quintana San Emeterio , L. Scodellaro , I. Vila , R. Villar Cortabitarte , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁷ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁵⁸ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , R. Ardino , E. Auffray , J. Baechler, D. Barney , J. Bendauid , M. Bianco , A. Bocchi , L. Borghonovi , C. Botta , A. Bragagnolo , C.E. Brown , C. Caillol , G. Cerminara , P. Connor , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , P.J. Fernández Manteca , W. Funk , A. Gaddi, S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, A. Gruber , J. Hegeman , J.K. Heikkilä , R. Hofsaess , B. Huber , V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁶ , G. Karathanasis , S. Laurila , P. Lecoq , C. Lourenço , A.-M. Lyon , M. Magherini , L. Malgeri , M. Mannelli , A. Mehta , F. Meijers , J.A. Merlin, S. Mersi , E. Meschi , M. Migliorini , F. Monti , F. Moortgat , M. Mulders , M. Musich , I. Neutelings , S. Orfanelli, F. Pantaleo , M. Pari , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabadý , A. Reimers , B. Ribeiro Lopes , F. Riti , P. Rosado , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁵⁹ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , E. Vernazza , J. Wanczyk⁶⁰ , J. Wang, S. Wuchterl , M. Zarucki , P. Zehetner , P. Zejdl , G. Zevi Della Porta 

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶¹ , L. Caminada⁶¹ , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , U. Langenegger , L. Noehte⁶¹ , T. Rohe , A. Samalan 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'archiacchiac⁶⁰ , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte , W. Lustermann , M. Malucchi , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani⁶⁰ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , B. Ristic , R. Seidita , J. Steggemann⁶⁰ , A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler⁶² , P. Bärttschi , F. Bilandzija , M.F. Canelli , G. Celotto , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , T.H. Kwok , S. Leontsinis , V. Lukashenko , A. Macchiolo , F. Meng , M. Missiroli , J. Motta , P. Robmann, M. Senger , E. Shokr , F. Stäger , R. Tramontano , P. Viscone 

National Central University, Chung-Li, Taiwan

D. Bhowmik, C.M. Kuo, P.K. Rout , S. Taj , P.C. Tiwari³⁷ 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu , E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Dolek , I. Dumanoglu⁶³ , Y. Guler⁶⁴ , E. Gurpinar Guler⁶⁴ , C. Isik ,
O. Kara , A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir⁶⁵ , B. Tali⁶⁶ ,
U.G. Tok , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁶⁷ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi⁶⁸ , E. Gülmez , M. Kaya⁶⁹ , O. Kaya⁷⁰ , M.A. Sarkisla⁷¹,
S. Tekten⁷² 

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{63,73} , S. Sen⁷⁴ 

Istanbul University, Istanbul, Turkey

O. Aydilek⁷⁵ , B. Haciasahinoglu , I. Hos⁷⁶ , B. Kaynak , S. Ozkorucuklu , O. Potok ,
H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci , C. Dozen⁷⁷ , B. Isildak⁷⁸ , E. Simsek , D. Sunar Cerci , T. Yetkin⁷⁷ 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , O. Dadazhanova, B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , D. Dharmender,
H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Parames-
varan , L. Robertshaw, M.S. Sanjrani⁴¹, J. Segal, V.J. Smith 

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁹ , C. Brew , R.M. Brown , D.J.A. Cockerill ,
A. Elliot , K.V. Ellis, J. Gajownik , K. Harder , S. Harper , J. Linacre , K. Manolopoulos,
M. Moallemi , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu ,
G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen ,
T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya ,
D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer,
G. Fedi , G. Hall , H.R. Hoorani , A. Howard, G. Iles , C.R. Knight , P. Krueper ,
J. Langford , K.H. Law , J. León Holgado , E. Leutgeb , L. Lyons , A.-M. Magnan ,
B. Maier , S. Mallios, A. Mastronikolis , M. Mieskolainen , J. Nash⁸⁰ , M. Pesaresi ,
P.B. Pradeep , B.C. Radburn-Smith , A. Richards, A. Rose , L. Russell , K. Savva ,
C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee²⁸ , M. Vojinovic ,
N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

Bethel University, St. Paul, Minnesota, USA

J.M. Hogan⁸¹ 

Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , S. Raj , A.E. Simsek , S.S. Yu 

The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , S. Campbell, R. Chudasama , S.I. Cooper , C. Crovella , G. Fidalgo , S.V. Gleyzer , A. Khukhunaishvili , K. Matchev , E. Pearson, C.U. Perez , P. Rumerio⁸² , E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

S. Cholak , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , J. Fulcher , F. Golf , S. Jeon , J. O'Cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , E. Wurtz, A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA

G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , T. Kwon , L. Lambrecht , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , J. Roloff, T. Russell , S. Sagir⁸³ , X. Shen , M. Stamenkovic , N. Venkatasubramanian 

University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , E. Cannaert, M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , O. Kukral , G. Mocellin , S. Ostrom , I. Salazar Segovia, J.S. Tafoya Vargas , W. Wei , S. Yoo 

University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca , A. Nunez Del Prado , D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , B. D'Anzi , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , V. Krutelyov , R. Lee , J. Letts , H. Li, M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch, M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , A. Yagil , Z. Zhao

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , S. Carron Montero⁸⁴ , K. Downham , C. Grieco , M.M. Hussain, J. Incandela , M.W.K. Lai, A.J. Li , P. Masterson , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang 

California Institute of Technology, Pasadena, California, USA

A. Albert , S. Bhattacharya , A. Bornheim , O. Cerri, R. Kansal , J. Mao , H.B. New-

man , G. Reales Gutiérrez, T. Sievert, M. Spiropulu , J.R. Vlimant , R.A. Wynne , S. Xie 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M. Cremonesi, V. Dutta , E.Y. Ertorer , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , P. Meiring , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , S. Kwan , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , J. Dickinson , A. Duquette, J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives , J. Monroy , G. Niendorf , M. Oshiro , J.R. Patterson , A. Ryd , J. Thom , P. Wittich , R. Zou , L. Zygala 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸⁵ , C.E. Perez Lara , F. Ravera , A. Reinsvold Hall⁸⁶ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , V. Sulimov , Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Goff, R. Hashmi , R.S. Kim , T. Kolberg , G. Martinez , M. Mazza , H. Prosper , P.R. Prova, R. Yohay 

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , M. Hohlmann , M. Lavinsky, E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , N. Barnett, A. Baty , C. Bennett , R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , C. Mills , S. Nanda , G. Nigmatkulov , B. Ozek , T. Phan, D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , J. Yoo 

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend , K. Dilsiz⁸⁷ , O.K. Köseyan , A. Mestvirishvili⁸⁸ , O. Neogi, H. Ogul⁸⁹ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁹⁰ 

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , S. Sekhar , M.V. Srivastav , M. Swartz 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁹¹ , C. Rogan , C. Royon , S. Rudrabhatla , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, USA

B. Allmond , N. Islam , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , G.G. Reddy , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim , S. Lascio , P. Major , A.C. Mignerey , C. Palmer , C. Papageorgakis , M.M. Paranjpe , E. Popova⁹² , A. Shevelev , L. Zhang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , H. Bossi , S. Bright-Thonney , I.A. Cali , Y.c. Chen , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov , G. Grosso , P. Harris , D. Hoang , G.M. Innocenti , D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , T.a. Sheng , G.S.F. Stephans , D. Walter , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , W.J. Jackson , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , R. Rusack , O. Sancar , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , A. Mandal , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar , A. Williams , P. Young , D. Yu 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma , M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , M. Lu , N. Manganelli , R. McCarthy , D.M. Morse , T. Orimoto , L. Skinnari , C.S. Thoreson , E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliencio , M. Velasco , J. Wang 

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci , S. Castells , A. Das , A. Ehnis , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , A. Karneyeu , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²² , H. Nelson , M. Osherson 

A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , R. De Los Santos , L.S. Durkin , C. Hill , M. Joyce ,
M. Nunez Ornelas , D.A. Wenzl, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , M. Galli ,
B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , Y. Lai , D. Lange , A. Loeliger ,
D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma 

Purdue University, West Lafayette, Indiana, USA

S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev ,
M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , N.R. Saha ,
J.F. Schulte , F. Wang , A. Wildridge , W. Xie , Y. Yao , Y. Zhong 

Purdue University Northwest, Hammond, Indiana, USA

N. Parashar , A. Pathak , E. Shumka 

Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , C. Arbour , T. Carnahan , K.M. Ecklund , S. Freed,
P. Gardner, F.J.M. Geurts , T. Huang , I. Krommydas , N. Lewis, W. Li , J. Lin ,
O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , M. Wulansatiti , E. Yigitbasi ,
Y. Zhang 

University of Rochester, Rochester, New York, USA

O. Bessidskaia Bylund, A. Bodek , P. de Barbaro[†] , R. Demina , A. Garcia-Bellido ,
H.S. Hare , O. Hindrichs , N. Parmar , P. Parygin⁹² , H. Seo , R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , S. Donnelly, D. Gadkari , Y. Gershtein ,
E. Halkiadakis , C. Houghton , D. Jaroslawski , A. Kobert , S. Konstantinou ,
I. Laflotte , A. Lath , J. Martins , M. Perez Prada , B. Rand , J. Reichert , P. Saha ,
S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , J. Harris, T. Holmes , A.R. Kanuganti ,
N. Karunarathna , J. Lawless, L. Lee , E. Nibigira , B. Skipworth, S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov , A. Bolshov, O. Bouhali⁹³ ,
A. Cagnotta , V. D'Amante , R. Eusebi , P. Flanagan , J. Gilmore , Y. Guo, T. Kamon ,
S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane ,
S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen ,
J. Zhang

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung , B. Cox , J. Hakala , G. Hamilton Ilha Machado, R. Hirosky , M. Jose, A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez , Z. Wu

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , S. Lomte , R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, T. Nelson, G. Parida , L. Pétré , D. Pinna , A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , R. Chistov⁹⁴ , M. Danilov⁹⁴ , T. Dimova⁹⁴ , A. Ershov⁹⁴ , S. Gninenko , I. Gorbunov , A. Gribushin⁹⁴ , A. Kamenev , V. Karjavine , M. Kirsanov , V. Klyukhin⁹⁴ , O. Kodolova⁹⁵ , V. Korenkov , I. Korsakov, A. Kozyrev⁹⁴ , N. Krasnikov , A. Lanev , A. Malakhov , V. Matveev⁹⁴ , A. Nikitenko^{96,95} , V. Palichik , V. Perelygin , S. Petrushanko⁹⁴ , S. Polikarpov⁹⁴ , O. Radchenko⁹⁴ , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹⁴ , K. Slizhevskiy, V. Smirnov , O. Teryaev , I. Tlisova⁹⁴ , A. Toropin , N. Voytishin , A. Zarubin , I. Zhizhin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

E. Boos , V. Bunichev , M. Dubinin⁸⁵ , V. Savrin , A. Snigirev , L. Dudko , K. Ivanov , V. Kim²² , V. Murzin , V. Oreshkin , D. Sosnov 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at The University of the State of Amazonas, Manaus, Brazil

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at China Center of Advanced Science and Technology, Beijing, China

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Also at School of Physics, Zhengzhou University, Zhengzhou, China

¹³Now at Henan Normal University, Xinxiang, China

¹⁴Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁵Now at The University of Iowa, Iowa City, Iowa, USA

¹⁶Also at Center for High Energy Physics, Peking University, Beijing, China

¹⁷Also at Cairo University, Cairo, Egypt

¹⁸Also at Suez University, Suez, Egypt

¹⁹Now at British University in Egypt, Cairo, Egypt

²⁰Also at Purdue University, West Lafayette, Indiana, USA

²¹Also at Université de Haute Alsace, Mulhouse, France

²²Also at an institute formerly covered by a cooperation agreement with CERN

- ²³Also at University of Hamburg, Hamburg, Germany
- ²⁴Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ²⁵Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ²⁶Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁷Also at Forschungszentrum Jülich, Jülich, Germany
- ²⁸Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁹Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary
- ³⁰Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- ³¹Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ³²Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ³³Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³⁴Also at The University of Kansas, Lawrence, Kansas, USA
- ³⁵Also at Punjab Agricultural University, Ludhiana, India
- ³⁶Also at University of Hyderabad, Hyderabad, India
- ³⁷Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁸Also at University of Visva-Bharati, Santiniketan, India
- ³⁹Also at IIT Bhubaneswar, Bhubaneswar, India
- ⁴⁰Also at Institute of Physics, Bhubaneswar, India
- ⁴¹Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴²Also at Isfahan University of Technology, Isfahan, Iran
- ⁴³Also at Sharif University of Technology, Tehran, Iran
- ⁴⁴Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁵Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁶Also at Helwan University, Cairo, Egypt
- ⁴⁷Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁸Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁹Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵⁰Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵¹Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵²Also at Lulea University of Technology, Lulea, Sweden
- ⁵³Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁴Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁵⁵Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁶Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁷Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁸Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁵⁹Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶⁰Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶¹Also at Universität Zürich, Zurich, Switzerland
- ⁶²Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶³Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁴Also at Konya Technical University, Konya, Turkey
- ⁶⁵Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁶Also at Adiyaman University, Adiyaman, Turkey

-
- ⁶⁷Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁶⁸Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey
- ⁶⁹Also at Marmara University, Istanbul, Turkey
- ⁷⁰Also at Milli Savunma University, Istanbul, Turkey
- ⁷¹Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
- ⁷²Also at Kafkas University, Kars, Turkey
- ⁷³Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁴Also at Hacettepe University, Ankara, Turkey
- ⁷⁵Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁶Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁷Also at Istinye University, Istanbul, Turkey
- ⁷⁸Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁹Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁸⁰Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸¹Also at Bethel University, St. Paul, Minnesota, USA
- ⁸²Also at Università di Torino, Torino, Italy
- ⁸³Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸⁴Also at California Lutheran University, Thousand Oaks, California, USA
- ⁸⁵Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁶Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸⁷Also at Bingöl University, Bingöl, Turkey
- ⁸⁸Also at Georgian Technical University, Tbilisi, Georgia
- ⁸⁹Also at Sinop University, Sinop, Turkey
- ⁹⁰Also at Erciyes University, Kayseri, Turkey
- ⁹¹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹²Now at another institute formerly covered by a cooperation agreement with CERN
- ⁹³Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
- ⁹⁴Also at another institute formerly covered by a cooperation agreement with CERN
- ⁹⁵Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁹⁶Also at Imperial College, London, United Kingdom